

This Provisional PDF corresponds to the article as it appeared upon acceptance. Fully formatted PDF and full text (HTML) versions will be made available soon.

A forgotten Collection: the Libyan Ethnobotanical exhibits (1912-14) by A. Trotter at the Museum O. Comes at the University Federico II of Naples, Italy

Journal of Ethnobiology and Ethnomedicine 2012, **8**:4 doi:10.1186/1746-4269-8-4

Antonino De Natale (antonino.denatale@gmail.com)
Antonino Pollio (anpollio@unina.it)

ISSN 1746-4269

Article type Research

Submission date 10 November 2011

Acceptance date 21 January 2012

Publication date 21 January 2012

Article URL <http://www.ethnobiomed.com/content/8/1/4>

This peer-reviewed article was published immediately upon acceptance. It can be downloaded, printed and distributed freely for any purposes (see copyright notice below).

Articles in *Journal of Ethnobiology and Ethnomedicine* are listed in PubMed and archived at PubMed Central.

For information about publishing your research in *Journal of Ethnobiology and Ethnomedicine* or any BioMed Central journal, go to

<http://www.ethnobiomed.com/authors/instructions/>

For information about other BioMed Central publications go to

<http://www.biomedcentral.com/>

A forgotten collection: the Libyan Ethnobotanical Exhibits (1912-14) by A. Trotter at the Museum O. Comes at the University Federico II in Naples, Italy

Antonino De Natale^{1*} and Antonino Pollio²

^{1*}Department of Soil, Plant, Environmental and Animal Production Sciences, University of Naples "Federico II" - Via Università, 100, 80055 Portici (NA), Italy.

² Department of Biological Sciences / Section of Plant Biology, University of Naples "Federico II" - Via Foria, 223, 80139 Napoli, Italy

*Corresponding author

Email

ADN: denatale@unina.it

AP: anpollio@unina.it

Abstract

Background - The Ethnobotanical Collection from the Libyan territories of the botanist Alessandro Trotter is included in the Oratio Comes Botanical Museum at the Faculty of Agraria at the University Federico II in Naples. Trotter explored different territories of Libya, mainly Tripolitania, between 1912-1924, collecting plant specimens and the drugs most frequently sold in the markets. The Libyan herbarium currently includes over 2300 sheets of mounted and accessioned plants. The drugs, mostly acquired by Trotter from Tripolitanian markets, were identified and packed in 87 paper sheets or boxes. Trotter added ethnobotanical information for each species when available.

Methods - A database of the herbarium species and the drugs has been carried out, after a taxonomic update. Nomenclature has been revised according to the African flowering plants database and the World Checklist of selected plant families, and a comparison with currently available ethnopharmacological data from North African has been attempted.

Results - In this study, ethnopharmacological data related to about 80 species of flowering plants and to 4 lichens are presented. The plants are mainly from Mediterranean or Sub-Saharan habitats and belong to 37 different families; Lamiaceae was the most cited family, with 10 accessions. Generally, the aerial parts of the plants are the most frequently used (28 species), followed by leaves (15 species), flowers and seeds (9 species), fruits (7 species) and hypogean organs (roots, rhizomes, tubers: 5 species). Plants were generally processed in very simple ways: infusion or decoction of the plants were prepared and orally administered or used for topical applications. A wide range of conditions was treated, ranging from mental disorders to skin affections. All the organs of human body are considered, but the pathologies of gastro-intestinal tract, respiratory system and those related to traumatic accidents were the most frequently mentioned. The comparison with the recent ethnopharmacological research in Maghreb and its neighboring countries reveals a high correspondence; almost all the plants cited by Trotter are still used in the folk medicine of at least one of the North African countries, and the therapeutic uses of each plant appear consistent over time.

Conclusions - The information collected by Trotter is an important contribution to tracing plant utilization in Libyan folk medicine over the last century.

Key words: Herbarium, North Africa, Libya, Traditional plant use, Medicinal plants

Background

The Orazio Comes Botanical Museum at the Facoltà di Agraria dell'Università Federico II di Napoli encompasses the Historical Library, the Herbarium Porticense (PORUN), the Mycological Herbarium (POR) and wood collections (Xilotomoteque and Xiloteque). These wide-ranging collections were created in the course of the 19th and 20th Centuries and were kept in different sections of the Faculty of Agraria. Only recently was the Polo Museale delle Scienze Agrarie instituted, with the aim of unifying all the collections under a common structure. While the definitive building is under re-construction, the collections of the O. Comes Museum have been temporarily transferred to a provisional site. This on-going reorganization provided an opportunity for a careful examination of the items from each collection, including the archives, which document the work of various botanists from the early 1600s to 1940 [1,2,3]. One of the most interesting results of this work was the rediscovering of the Botanical Collection from the Libyan territories of the botanist Alessandro Trotter (**Figure 1**). A. Trotter was born on 1874 in Udine, Italy. In 1899, he graduated with a degree in Botany from the University of Padua, under Professor A. Saccardo, who accepted him as his assistant immediately after his graduation. A. Trotter remained at the University of Padua until 1902, when he was entrusted to teach Plant Pathology at the School of Enology of Avellino, and, subsequently, at the Agronomy Faculty of Portici, Naples, where he worked until 1949 as full professor of Botany. Trotter's scientific career reflected his broad interest in Botany. Since his beginning at the University of Padua, he demonstrated a strong attitude towards the floristic, with a specific interest in mycology, but he was also committed to the study of horticultural species, particularly those from South Italia. On behalf of the Italian Government, from 1912 to 1924, Trotter participated in different expeditions in Libya [4] (**Figure 2**), mainly in Tripolitania [5,6,7]. The results of Trotter's work in this region and, to a lesser extent, in other Libyan territories was a collection of plant specimens and a *repertorium* of the drugs he bought in the markets of Tripolitania, together with an archive of information, field notes, and photographs. Some of his findings were published in two reports [8,9], which are unavailable to the current scientific community. Over time, Trotter's materials were scattered throughout different

rooms of the Botany Department and merged with other accessions belonging to different collections. The organization of the new museum offered the possibility of grouping Trotter's exhibits in one collection, which represents a unique source of ethnobotanical information, dating back to a century ago, on a scarcely known geographical area of North Africa. It has been stressed that there are only a few places in the world for which diachronic data for ethnopharmacological purposes are available [10]. Recent efforts on this subject have been recently presented for Estonia [11] and Poland [12,13], but old ethnobotanical data are very scarce for the African continent [14,15].

The aim of this paper is to report the data on plants used in folk medicine of Tripolitania and neighboring regions collected by Trotter during his expeditions in Libya and to present a preliminary description of the exhibits now available in the Libyan ethnobotanical collection of the Orazio Comes Museum.

Methods

Geographical description of the study area

North Tripolitania is dominated by the Jaffar Plain, which slopes downward to the west and upward to the highland territories (Jabal Nafusa), which in turn continue eastward with a belt of hills around Tarhuna and Homa [16]. The vegetation is dominated by steppes, which cover a zone extending from the coast west of Zamiya to a large part of Jaffara Plain, the Jabal Nafusa, and eastward to the Misurata region [17]. The soils around Tripoli, extending to central Jaffara, are rich in organic compounds, but the soils are poor and sandy in the rest of North Tripolitania and rocky in Jabal Nafusa. In coastal Tripolitania and in the Jabal Nafusa highlands, the climate is Mediterranean, with long, dry and hot summers, whereas in central and eastern Jaffara and in inland Jabal, the climate is semi arid, and drought occurs frequently [18]. At the beginning of 20th century, agriculture was concentrated in the oasis of the coastal territories and in part of the Jabal Nafusa plain. In the rest of the region, pastoralism (sheep, goats and camels) was the most common land use, together with transhumant cereal cultivation. The areas of mixed cereal cultivation and pastoralism were held by seminomadic groups through a combination of private and collective tenure [16].

History of the Trotter Libyan Collections

After his first expedition to Libya (1912), Trotter assembled a collection of plants primarily from North Tripolitania. In the subsequent trips, he gathered specimens from other regions of Libya as well [5,6,7].

The Trotter Collection, now located in the Orazio Comes Museum, features herbarium samples, drugs, historical photographs, and manuscripts.

The Libyan Phanerogamic collection currently includes over 2300 sheets of mounted and accessioned plants. The herbarium is organized alphabetically by family, and each herbarium sheet measures 43 x 28 centimeters (**Figure 3**). Paper specimen labels are attached to the right corner of the herbarium sheet. Loose parts or fragments of plant organs are included in envelopes attached to the herbarium sheet. Each voucher specimen was annotated by Trotter with the current taxonomic name at the time, the Libyan location from which the specimen was acquired, and the collection date. Additionally, Trotter added ethnobotanical information for about 80 species, which have now been grouped in a specifically designed archive.

The drugs (**Figure 4**), prevalently acquired by Trotter from Tripolitania markets, were identified and packed by Trotter in 83 paper sheets or boxes (**Figure 5**), which were stored in a wooden cupboard (**Figure 6**). Included in the collection were some unidentified drugs, corresponding to n. 74 and 83.

Recently, a database of the herbarium species and drugs has been carried out, and, subsequently, a taxonomic update has been attempted. The nomenclature has been revised according to the African flowering plants database [19] and the World Checklist of selected plant families [20].

Results

Italian colonization of Tripolitania began in 1911 without a preliminary assessment of the land: for this reason, geographical expeditions in Libyan territories were undertaken between 1912 and 1916, with the participation of various specialists. The first expedition [4] had the task of carrying out mineralogical and agronomic studies on Tripolitania, and Trotter was the agronomist of the expedition. During his residence in Libya, he studied the flora (algae, fungi, and higher plants, including cultivated species) of Tripolitania. He was also greatly involved in the collection of

information on plant uses from the local people. This interest was driven by practical needs, as plants used by Libyans could have potential economic benefits to emigrant settlers. In this paper, we have concentrated our attention on the medicinal uses of Libyan plants collected by Trotter. In Table 1 [21-45], these ethnopharmacological data are compared with the recent available information from Libya and other neighboring countries.

Trotter collected his ethnobotanical data mainly during the first expedition (1912). The provenance of the information is not always indicated, but we are far from the present standards of ethnobotanical investigations. Frequently, a single or very few informants are the source of the data, and, in some cases, Trotter reported only that the plant was used as a medicine, without describing the therapeutic applications. Notwithstanding these limitations, a not-irrelevant body of knowledge was assembled.

Data are related to about 70 species of flowering plants and to 4 lichens. The plants are mainly from Mediterranean or Sub-Saharan habitats, with a slight prevalence for the former. The plants belong to 37 different families; Lamiaceae was the most cited family with 10 accessions, followed by Asteraceae with 9. The prevalence of these families has been reported in the conclusive report of the Rubia project on medicinal plants in Mediterranean area [46]. The authors suggest that this could be because the plants belonging to these families are easily recognizable by the morphology of the flowers and due to their aromas and flavors.

Generally, the aerial parts of the plants are the most frequently used (28 species), followed by leaves (15 species), flowers and seeds (9 species), fruits (7 species) and hypogean organs (roots, rhizomes, tubers: 5 species).

Plants were generally processed in very simple ways: an infusion or decoction of the plants were prepared and orally administered or used for topical applications. A wide range of affections was treated, ranging from mental disorders to skin affections. All the organs of human body were considered, but the pathologies of the gastro-intestinal tract, respiratory system and those related to traumatic accidents (contusions, swellings, burnings, wounds) were the most frequently mentioned.

Trotter described a decoction, named *sciba*, made with three different lichens belonging to the genera *Evernia* and *Ramalina*. The species mentioned are found

worldwide and largely used for dyeing [47], but the species are also used in perfumery [48] and in traditional medicine of numerous countries, due to the presence of active compounds, as usnic acid and atranorin [49]. Trotter did not indicate the therapeutic applications for this decoction. With a similar name, al-sheba, the lichen *Parmotrema tinctorium* is sold as food spice in India [50]. As far we know, there are no other reports dealing with this preparation in the Maghreb countries.

Overall, 75 plants (including the four lichens previously mentioned) were also collected as drugs by Trotter; 30 of these had medicinal properties. Two mineral samples (antimony and clay) were also held in the drug collection (**Figure 7** and **Figure 8**); they were used in topical wound treatment by numerous ancient and primitive societies [51]. The drugs were mainly sold in the Tripoli market, but also in local markets spread along Tripolitania. Libya is located in the middle of Mediterranean and was an important crossroad for trade in ancient times. The Libyan towns established commercial relationships with countries of all three continents, Africa, Asia and Europe. The town of Cyrene was a prime center for the export of the medicinal herb called silphium, one of the essential commodities of the Mediterranean region in classical era [52]. A late echo of the ancient flourishing trades was still present in the drugs found by Trotter at the beginning of the 20th century. He collected drugs that are, to a large extent, of Mediterranean origin, and are currently traded in Mediterranean region. Drugs from herbs such as *Ajuga iva* or *Artemisia arborescens* or those belonging to the genera of Lamiaceae, listed in Table 1, are commonly found in the markets along the Mediterranean, from Moroccan bazaars [44] to the herbal shops of Turkey [40] and Greece [53]. In contrast, some drugs were of Asiatic provenance, such as *Alpinia officinarum* [54] and *Piper retrofractum* [55], whereas others, such as *Tanacetum parthenium*, came mainly from Europe [56], suggesting that the ancient trade routes from Asia and Europe to North Africa were still being used a century ago. Few drugs were produced from indigenous plants; perhaps the most interesting case is that of *Ferula marmarica*, a plant native to some Libyan regions [34], which was used in classical times to produce the ammoniac gum [35], to which Dioscorides attributed relevant therapeutic roles ranging from anti-inflammatory to digestive and painkiller [57].

Conclusion

The Trotter Collection can represent a useful tool for current ethnopharmacological research in Libya and neighboring countries. It is known that about 80% of the African population presently rely on traditional forms of health care, but it is not easy to document continuity and changes in therapeutic approaches. The information collected by Trotter contributes to filling this gap, enabling us to trace the use of plant utilization in Libyan folk medicine over the last century. A comparison with the recent ethnopharmacological research in Maghreb reveals a high correspondence; almost all of the plants cited by Trotter are still used in the folk medicine of at least one of the North African countries, and therapeutic uses of each plant appear consistent over that time.

The results of this study, although based on information that needs to be confirmed by current methodologies, seem to suggest that it is possible to find a core of a shared popular medicine along the African Coast of the Mediterranean Sea, probably due to climatic, cultural and linguistic continuity.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

The authors contributed equally to this paper. All authors read and approved the final manuscript.

References

1. De Natale A: **Herbarium Porticense**. In *I Musei delle Scienze Agrarie. L'evoluzione delle Wunderkammern*. (Eds. S Mazzoleni and S Pignattelli). Univ. degli Studi di Napoli "Federico II", Napoli; 2007, pp. 52-74.
2. De Natale A, Cellinese N: **Imperato, Cirillo, and a series of unfortunate events: a novel approach to assess the unknown provenience of historical herbarium specimens**. *Taxon* 2009, **58(3)**:963-970.
3. De Natale A, Marziano F: **Note di etnomicologia per il territorio di Catania (Sicilia), in un manoscritto inedito di Fridiano Cavara**. *Micologia Italiana* 2009, **38(3)**:51-60.
4. Galli P: **La Tripolitania. Carta Geografica Dimostrativa**. Antonio Vallardi Ed., Milano; [s.d.] (about 1912).
5. De Cillis E, Franchi S, Trotter A, Tucci F: **Ricerche e studi agrolgici sulla Libia**. Vol. 1. La zona di Tripoli. Ministero di Agricoltura, Industria e Commercio, Istituto Italiano d'Arti Grafiche (Bergamo); 1912.
6. Trotter A: **A proposito di alcune piante del Gébel tripolitano**. *Bullettino dell'Orto Botanico della Regia Università di Napoli* 1914, **4**:235-238.
7. Trotter A: **Osservazioni botaniche in Tripolitania**. *Rivista della Tripolitania* 1924, **1(3)**:1-24.
8. Trotter A: **Nuove osservazioni botaniche sulla Tripolitania**. In *La Tripolitania Settentrionale*. Vol. 1 (Relazione della Commissione Ministeriale per lo studio agrolgico della Tripolitania). Tip. Naz. Bertero; 1913.
9. Trotter A: **Flora economica della Libia**. Tipografia dell'Unione Editrice, Roma; 1915.
10. Heinrich M, Kufer J, Leonti M, Pardo-de-Santayana M: **Ethnobotany and ethnopharmacology-interdisciplinary links with the historical sciences**. *Journal of Ethnopharmacology* 2006, **107**:157-160.
11. Sõukand R, Kalle R: **Change in medical plant use in Estonian ethnomedicine: A historical comparison between 1888 and 1994**. *Journal of Ethnopharmacology* 2011, **135**:251-260.
12. Łuczaj Ł: **Change in the utilization of wild green vegetables in Poland since the 19th century: A comparison of four ethnobotanical survey**. *Journal of Ethnopharmacology* 2010, **128**:395-404.
13. Łuczaj Ł: **Change in assumption day herbal bouquets in Poland: a nineteenth century study revisited**. *Economic Botany* 2011, **65(1)**:66-75.
14. Fossi Innamorati T: **Notizie di medicina popolare africana nell'Erbario Tropicale di Firenze**. *Webbia* 1973, **28(1)**:81-133.
15. Sandberg F, Perera-Ivarsson P, Rushdey El-Seedi H: **A Swedish collection of medicinal plants from Cameroon**. *Journal of Ethnopharmacology* 2005, **102**:336-343.
16. Fowler GL: **Italian colonization of Tripolitania**. *Annals of Association of American Geographers* 1972, **62(4)**:627-640.
17. Principi P: **I principali tipi di suolo della Tripolitania**. *Rivista di Agricoltura Subtropicale e Tropicale* 1947, **41**:195-201.
18. Fantoli A: **La siccità in Libia**. *Reale Accademia dei Lincei* 1935, **4(2)**:1-92.
19. Conservatoire et Jardin botaniques de la Ville de Genève e South African National Biodiversity Institute: **African Flowering Plants Database**. (version 3.2, last update 01.12.2009) Pretoria. [<http://www.ville-ge.ch/musinfo/bd/cjb/africa/>]
20. WCSP, **World Checklist of Selected Plant Families**. Facilitated by the Royal Botanic Gardens, Kew. [<http://apps.kew.org/wcsp/home.do>] (accessdate 2011-04-01)
21. IUCN, The World Conservation Union Centre for Mediterranean Cooperation Parque Tecnológico de Andalucía: **Medicinal Plants of North Africa**. [<http://data.iucn.org/places/medoffice/nabp/database/>] (last update 2005)
22. Le Houérou HN: **Plant diversity in Marmarica (Libya & Egypt): a catalogue of the vascular plants reported with their biology, distribution, frequency, usage, economic potential, habitat and main ecological features, with an extensive bibliography**. *Candollea* 2004, **59(2)**:259-282.
23. Boulos L: **Medicinal plants of North Africa**. References Publications Inc.; 1983.
24. Merzouki A, Ed-derfoufi F, Molero Mesa J: **Contribution to the knowledge of Rifian traditional medicine. II: Folk medicine in Ksar Lakbir district (NW Morocco)**. *Fitoterapia* 2000, **71**:278-307.

25. Bellakhdar J, Claisse R, Fleurentin J, Younos C: **Repertory of standard herbal drugs in the Moroccan pharmacopoea.** *Journal of Ethnopharmacology* 1991, **35**:123-143.
26. El-Hilaly J, Hmammouchi M, Lyoussi B: **Ethnobotanical studies and economic evaluation of medicinal plants in Taounate province (Northern Morocco).** *Journal of Ethnopharmacology* 2003, **86**:149-158.
27. De Boer HJ: **African plants as antipathogen agents: efficacy and clinical evidence.** In *Botanical Medicine in clinical practice* (RR Watson and VR Preedy Eds.). CABI, Cromwell Press, Trowbridge, UK; 2008:13-19.
28. Hammiche V, Maiza K: **Traditional medicine in Central Sahara: Pharmacopoeia of Tassili N'ajjer.** *Journal of Ethnopharmacology* 2006, **105**:358-367.
29. Maiza K, Brac De La Perrière RA, Hammiche V: **Pharmacopée traditionnelle saharienne: Sahara septentrional.** In *Actes du 2^e Colloque Européen d'Ethnopharmacologie et de la 11^e Conférence internationale d'Ethnomédecine, Heidelberg, 24-27 mars 1993, Heidelberg.* Medicines and Foods: The Ethnopharmacological Approach. Heilmittel und Nahrungsmittel aus ethnopharmakologischer Sicht, Heidelberg, AGEM; 1993:169-171.
30. El-Kamali HH, El-Kalifa KF: **Folk medicinal plants of riverside forest of the Southern Blue Nile district, Sudan.** *Fitoterapia* 1999, **70**:493-497.
31. Hussein Ayoub SM, Baerheim Svendsen A: **Medicinal and aromatic plants in the Sudan: usage and exploration.** *Fitoterapia* 1981, **52**:243-246.
32. Basar S: **Phytochemical investigation on Boswellia species.** PhD thesis, University of Hamburg; 2005.
33. Hussein Kotb FT: **Medicinal plants in Libia.** Arab Encyclopedia House, Beirut-Lebanon; 1985.
34. El-Darier SM, El-Mogaspi FM: **Ethnobotany and relative importance of some endemic plant species at El-Jabal El-Akhdar region (Lybia).** *World Journal of Agricultural Sciences* 2009, **5(3)**:353-360.
35. Stapf O: **The gums ammoniac of Morocco and the Cyrenaica.** *Bullettin of Miscellaneous Information. Royal Botanic Garden, Kew* 1907, **10**:375-388.
36. Ghirardini MP, Carli M, del Vecchio N, Rovati A, Cova O, Valigi F, Agnetti G, Macconi M, Adamo D, Traina M, Laudini F, Marcheselli I, Caruso N, Gedda T, Donati F, Marzadro A, Russi P, Spaggiari C, Bianco M, Binda R, Barattiere E, Tognacci A, Girardo M, Vaschetti L, Caprino P, Sesti E, Andreozzi G, Coletto E, Belzer G, Pieroni A: **The importance of a taste. A comparative study on wild food plant consumption in twenty-one local communities in Italy.** *Journal of Ethnobiology and Ethnomedicine* 2007, **3**:22. [<http://www.ethnobiomed.com/content/3/1/22>]
37. Nebel S, Heinrich M: **Ta Chòrta: A Comparative Ethnobotanical-Linguistic Study of Wild Food Plants in a Graecanic Area in Calabria, Southern Italy.** *Economic Botany* 2009, **63(1)**:78-92.
38. Botsaris SA: **Plants used traditionally to treat malaria in Brazil: the archives of Flora Medicinal.** *Journal of Ethnobiology and Ethnomedicine* 2007, **3**:18. [<http://www.ethnobiomed.com/content/3/1/18>]
39. Lev E, Amar Z: **"Fossils" of practical medical knowledge from medieval Cairo.** *Journal of Ethnopharmacology* 2008, **119**:24-40.
40. Everest A, Ozturk E: **Focusing on the ethnobotanical uses of plants in Mersin and Adana provinces (Turkey).** *Journal of Ethnobiology and Ethnomedicine* 2005, **1**:6. [<http://www.ethnobiomed.com/content/1/1/6/>]
41. Hamdan II, Afif FU: **Studies on the in vitro and in vivo hypoglycemic activities of some medicinal plants used in treatment of diabetes in Jordanian traditional medicine.** *Journal of Ethnopharmacology* 2004, **93(1)**:117-121.
42. Isman MB: **Botanical insecticides, deterrents, in modern agriculture and an increasingly regulated world.** *Annual Reviews Entomology* 2006, **51**:45-66.
43. Lahsissene H, Kahouadji A, Tijane M, Hseini S: **Catalogue des plantes medicinales utilisees dans la region de Zaër (Maroc Occidental).** *Lejeunia* 2009, n.s. **186**:1-25.
44. Fogg W: **The wares of Moroccan folk-doctor.** *Folklore* 1941, **52(4)**:273-303.
45. Setty AR, Sigal AH: **Herbal medications commonly used in the practice of rheumatology: Mechanisms of action, efficacy, and side effects.** *Semin Arthritis Rheum* 2005, **34**:773-784.
46. González-Tejero MR, Casares-Porcel M, Sánchez-Rojas CP, Ramiro-Gutiérrez JM, Molero-Mesa J, Pieroni A, Giusti ME, Censorii E, de Pasquale C, Della A, Paraskeva-Hadijchambi D, Hadjichambis A, Houmani Z, El-Demerdash M, El-Zayat M, Hmammouchi M, ElJohrig S: **Medicinal plants in the Mediterranean area: Synthesis of the results of the project Rubia.** *Journal of Ethnopharmacology* 2008, **116**:341-357.
47. Uphof CJTh: **Dictionary of economic plants.** J. Cramer Ed., New York; 1968.
48. Romagni JG, Dayan FE: **Structural diversity of lichen metabolites and their potential use.** In *Advances in microbial toxin research.* (RK Uphadhy Ed.). Kluwer Academic/Plenum Press, New York; 2002:151-158.
49. Malhotra S, Subban R, Singh A: **Lichens-Role in Traditional Medicine and Drug Discovery.** *The Internet Journal of Alternative Medicine* 2008, **5(2)**. [<http://www.ispub.com/ostia/index.php?xmlFilePath=journals/ijam/vol5n2/lichen.xml>]
50. Upreti DK, Divakar PK, Nayaka S: **Commercial and ethnic use of lichens in India.** *Economic Botany* 2005, **59(3)**:269-273.
51. Forrest RD: **Early history of wound treatment.** *Journal of the Royal Society of Medicine* 1982, **75**:198-205.

52. Roselli A: **Breve storia del silfio**. *Annali di Archeologia e storia antica* 2001, **8**:11-20.
53. Hanlidou E, Karousou R, Kleftoyanni V, Kokkini S: **The herbal market of Thessaloniki (N Greece) and its relation to the ethnobotanical tradition**. *Journal of Ethnopharmacology* 2004, **91(2-3)**:281-299.
54. Hare CP: **Indian medicinal plants: an illustrated dictionary**. Springer Verlag Berlin, Heidelberg; 2007.
55. Simonati CN, Helmy MA, Moise G: **Production of medicinal plants in Asia**. *Scientific Papers Management, Economic Engineering in Agriculture and Rural Development* 2009, **9(1)**:135-138.
56. Groenewegen AW, Knight DW, Heptinstall S: **Progress in medicinal chemistry of the herb feverfew**. In *Progress in Medicinal Chemistry* (GP Ellis and DK Luscombe Eds.). Elsevier 1991, **29**:217-238.
57. Lily Beck Y: **Dioscorides. De Materia Medica**. *Altertumswissenschaftliche Texte und Studien* 38. Hildesheim: Olms; 2005.

Figure legends

Figure 1 - A portrait of Alessandro Trotter.

Figure 2 - Geographical map of Libya in 1912 [4].

Figure 3 - Herbarium specimen from Trotter collection, *Artemisia herba-alba* Asso.

Figure 4 - A drug package.

Figure 5 - Paper sheets and boxes containing the drugs.

Figure 6 - The original wooden cabinet of Trotter Herbarium.

Figure 7 - Antimonium (PORUN - TTD76), Tripolitania Trotter collection Drug section.

Figure 8 - Clay (PORUN - TTD77), Tripolitania Trotter collection Drug section.

Table

Table 1 - Plants used in Lybian folk therapy at the beginning of XXth Century according to Trotter data.

Botanical name	Local names	Part(s) used	Description of uses	Present uses in Libya and neighboring Countries
<i>Aaronsohnia pubescens</i> (Desf.) K. Breimer et Humphries (PORUN - TTF2300) ASTERACEAE	<i>uás-uása</i> (arab); (tuaregh)	Leaves	Grounded and eaten against gastro-intestinal aches	The whole plant in infusion is used for gastrointestinal troubles and kidney stones. It is collected and sold in the markets of South Algeria [21]
<i>Achillea maritima</i> (L.) Ehrend. et Y.P. Guo (PORUN - TTF2318) ASTERACEAE	<i>agbita, sciba</i> (Algeria)	Aerial part	Sold as a febrifuge and emmenagogues	In Marmarica, it is known as medicinal plant [22]. In North African countries, the flowering branches are considered a febrifuge, emmenagogue, tonic, and taenifuge [23]
<i>Ajuga iva</i> (L.) Schreb. (PORUN - TTF2318) LAMIACEAE Additional file 1	<i>sciandagúra</i> (arab); <i>assaron?</i> (Cirenaica); <i>tletúl</i> (berber)	Aerial part	A cold infusion is anthelminthic; mixed with other components, against pulmonary affections	In Marmarica, the plant is considered medicinal [22]. A decoction of the aerial parts is administered against rheumatic pains, and as a carminative and stomachic. Aerial parts are also used as incense in ritual practices [24]. In Morocco, the hot infusion is considered antidiabetic [23]; also used as an anthelmintic and for intestinal disorders [25]. Crushed leaves and seeds are smoked for their narcotic effects [26]
<i>Albè vera</i> L. (PORUN - TTF2331) ASPHODELACEAE	<i>sabbàr</i>	Leaf juice	Medicinal properties (not described)	The transparent gel from leaf pulp is used as a vulnerary and laxative in different African countries [27]
<i>Alpinia officinarum</i> Hance (PORUN - TTF2331) ZINGIBERACEAE Additional file 2	<i>cúlgan, cúlgian</i>	Rhizome	Tonic	Drug imported from East Asia and used in Egypt and Morocco, used as an antitussive and stimulant [22,25]. A preparation from the root is used for rheumatism and sexual impotence [24]
<i>Arbutus unedo</i> L. (PORUN - TTF895) ERICACEAE	<i>sc'meri</i> (arab); <i>linz</i> (Algeria); <i>isisnu, sciscnu</i> (berber)	Bark of roots and leaves	Astringent	The decoction of leaves or raw fruits to treat kidney diseases [26]
<i>Artemisia arborescens</i> L. (PORUN - TTF895) ASTERACEAE Additional file 3	<i>sézeret Marian</i> (arab); <i>scih</i>	Young shoots, flowers, leaves	A decoction against intestinal affections	In Morocco, the plant is considered anthelmintic, aperitive, diuretic, emmenagogue and abortive [25]. A leaf infusion to treat common colds, vertigo, and helminthiasis. Powdered leaves are externally applied for skin infections and wrinkles [24]
<i>Artemisia campestris</i> L. subsp. <i>variabilis</i> (Ten.) Greuter (PORUN - TTF484) ASTERACEAE	<i>sc'ahâl</i> (arab); <i>togoft, tegoft,</i> <i>taghert, tághiat</i> (berber); <i>teghoch</i> (tuaregh)	Flowers, leaves	Anthelminthic, known as <i>duah lehnâsc</i>	In Algeria, it is used as a substitute for <i>A. absinthium</i> [23]. In Tassili N'ajjer (South Algeria), aerial parts and flowers are used for post-partum care, emmenagogue, analeptic, and antispasmodic. Also used as an anthelmintic, for stomach and liver affections. Vulnerary [28]

Artemisia herba-alba Asso (PORUN - TTF482) ASTERACEAE
Figure 3 -

scih (arab); *aghares* (berber); *azezzeré, zezeri* (temahac) and the inflorescence

In the Fezzan, the dried plant is drunk in North Sahara against digestive troubles, as an anthelmintic, and to treat eye affections and obesity [29]. A branch decoction is used for rheumatic pains and helminthiasis [24]. In Morocco, the plant is administered against gastrointestinal affection, as an antiseptic, anthelmintic, antidote to poisons, hypoglycemic and emmenagogue [25]

Roots, shoots and leaves are used from Egypt to Morocco as a tonic and stomachic, against headache, liver affections rheumatism, and for treatment of syphilis [23]

not reported

Leaves

Diuretic

Asphodelus refractus Boiss. (PORUN - TTF2304) ASPHODELACEAE

Balanites aegyptiaca (L.) Del. s.l. (PORUN - TTF2302) BALANITACEAE

hagilidi, heglig (arab); *tebôrac* (temahac); *tsciaisciot* (Tuat); *addua* (Hausa)

Young leaves, pulp of fruits

Leaves to clean sores; pulp fruit used in spleen diseases and to kill Guinea worms (*Vena medensis*)

The leaf decoction is used in the central Sahara (Algeria) and in other North African countries, as an anthelmintic, against skin affections, and as a vulnerary. Also administered to cure stomach, liver, pulmonary and spleen affections [23,28]. In Sudan, the fruit is administered as a purgative, a bark decoction against jaundice, and branch fumigation against rheumatism [30]. Also used as purgative and for bilharzias [31]

Boswellia sacra Flüeckiger (PORUN - TTF2305) BURSERACEAE
Additional file 4

Resin of stem

The yellowish granules of the resin are burnt and chewed for their aphrodisiac properties

not reported

Seeds

Medicinal properties (not described)

Brassica napus L. s.l. (PORUN - TTF2308) BRASSICACEAE

Capparis orientalis Veillard in Duh. (PORUN - TTD7; TTF101) CAPPARACEAE
Additional file 5

Leaves

Near Tripoli, a decoction of the plant is mixed with other herbs as a stomachic

léft (arab); *afrán* (temahac)

cábbar (arab); *tilult* (berber)

Seeds

Medicinal properties (not described)

Citrullus colocynthis (L.) Schrad. (PORUN - TTF2316) CUCURBITACEAE

handel, handla (arab); *taghillilit* (berber); *tassellet* (mezabita); *alched, hagi* (tuaregh)

Seeds

Seed infusion against the viper bites

Roots are used as an abortive. In Morocco, it is considered as an antiepileptic, anthelmintic, aphrodisiac, and hypoglycemic. Also administered against gonorrhoea and tinea [25]. Moreover, the fruit, broken into small pieces, is used to protect woolen clothing from moths [21]

In Marmarica region, it is known as fodder and as a medicinal plant [22]. A potion of the plant is drunk in North Sahara against digestive troubles, as an anthelmintic, and to treat eye affections and obesity [29]. A branch decoction is used for rheumatic pains and helminthiasis [24]. In Morocco, the plant is administered against gastrointestinal affection, as an antiseptic, anthelmintic, antidote to poisons, hypoglycemic and emmenagogue [25]

Roots, shoots and leaves are used from Egypt to Morocco as a tonic and stomachic, against headache, liver affections rheumatism, and for treatment of syphilis [23]

The leaf decoction is used in the central Sahara (Algeria) and in other North African countries, as an anthelmintic, against skin affections, and as a vulnerary. Also administered to cure stomach, liver, pulmonary and spleen affections [23,28]. In Sudan, the fruit is administered as a purgative, a bark decoction against jaundice, and branch fumigation against rheumatism [30]. Also used as purgative and for bilharzias [31]

The species grows in South Arabia and is known by the Arabians as "maghrayt d'sheehaz". The resin is traditionally sold in African markets as a disinfectant and also used in the preparation of cosmetics [32]

Included among Libyan medicinal plants as an emollient (roots) [33]

In North Sahara, it is administered as a pain-killer, mainly to treat toothaches, and against rheumatism [29]. In central Sahara (Tassili N'ajjer, Algeria), a poultice of fresh leaves is typically applied against rheumatism pains and headache [28]. Flowers and fruits are macerated to treat rheumatism [26]. An infusion of root bark is used as a cholagogue [24]

<i>Coriandrum sativum</i> L. (PORUN - TTF2313) APIACEAE	<i>cússbur, cóssbor</i> (arab)	Fruits	A syrup made from the fruit used against pulmonary affections	Aphrodisiac, anti-inflammatory, tonic. Seed and leaf decoction used against kidney stones, intestinal pains, insomnia, and scurvy [25,26]. Fruits are placed onto fire to remove bad spirits [24]
<i>Cuminum cyminum</i> L. (PORUN - TTD9) APIACEAE	<i>chemún ahdar</i> (arab); <i>azcar</i> (tuaregh)	Fruits	Chewed disorders	A milk infusion of fruit is administered against gastric pains and as an intestinal antiseptic, carminative and sudorific [24,25]. Also used in veterinary medicine [23]
Additional file 6 <i>Cupressus sempervirens</i> L. (PORUN - TTD60; TTF854) CUPRESSACEAE	<i>arz, sírua</i> (arab)	Seeds	An infusion against cough and heart diseases	In Libya, Morocco and Tunisia, the cones are considered antidiarrheal, antihemorrhagic, astringent, diuretic, expectorant, and sudorific [23]. In the El-Jabal El-Akhdar region (Libya), leaves and cones are administered in different ways against asthma, piles, and vaginal discharge [34]
Additional file 7				
<i>Curcuma longa</i> L. (PORUN - TTF2320) ZINGIBERACEAE	<i>córcob</i>	Roots	Medicinal properties (not described)	In Morocco, it is used as a digestive, stimulant, for blood diseases, and against amnesia [25]. The powdered rhizome is taken orally as a condiment, tonic, caefacient and digestive [24]
<i>Cymbopogon schoenanthus</i> (L.) Spreng. s.l. (PORUN - TTD79) POACEAE	<i>led'her</i> (Mizda, arab); <i>bu'rucúba, semmad</i> (Algeria); <i>lemmed, tiberrimt, leberint</i> (Algeria)	Basal part of the plant	Fumigation against a kind of influence called ahón	The plant is used in different countries of North Africa as an antihirheumatic, diuretic, emmenagogue, and febrifuge [23]
Additional file 8 <i>Cynodon dactylon</i> (L.) Pers. (PORUN - TTF1127) POACEAE	<i>négem</i> (arab); <i>tobbalt</i> (berber); <i>asezmir</i> (mezabita); <i>oscerar</i> (temahac)	Whole plant	The decoction is a diuretic	Diuretic [25]. Root decoction against stomach disorders [26]. A decoction of the rhizome or aerial part against rheumatism, kidney stones, uterine and menstrual pains, and abortion [24]
<i>Cyperus rotundus</i> L. s.l. (PORUN - TTF2307) CYPERACEAE	<i>giaâd</i> (arab)	Tubers	A water potion against heart diseases	North African Countries. The tubercles are used to cure a wide range of affections [23] In Sudan, a tuber decoction is used to treat stomach troubles and as an anthelmintic [30]. Stem galls mixed with leaves of <i>Lawsonia inermis</i> are powdered, kneaded with water and applied as a hair tonic [24]. The tubers are used to increase body weight [21]
<i>Elettaria cardamomum</i> (L.) Maton (PORUN - TTD8) ZINGIBERACEAE	<i>cacúla</i>	Fruits	Fruit is grounded and mixed with honey into a tonic	In Morocco, the plant is considered to be a cosmetic and aromatic and is used in hair care [25]
Additional file 9				
<i>Ephedra alata</i> Decne. s.l. (PORUN - TTF2310) EPHEDRACEAE	<i>alenda</i> (arab); <i>timatrat</i> (temahac)	Seeds and young shoots	Medicinal properties (not described)	Used in Algeria, Libya, and Morocco as an anthiasthmatic, anti-hypertensive, astringent, and depurative as well as used against headaches and for treatment of pulmonary affections [23]

<i>Euphorbia guyoniana</i> Boiss. et Reut. (PORUN - TTF918) <i>Euphorbia paralias</i> L. (PORUN - TTF922) EUPHORBIACEAE	<i>lebbin</i> (arab)	Latex	Topical application against viper bites	Different <i>Euphorbia</i> species are used in Morocco to treat skin diseases [26]
<i>Ferula marmarica</i> Asch. et Schweinf. (PORUN - TTD22) APIACEAE	<i>fassuch</i>	Gum resin	Medicinal properties described)	(not Source of gum ammoniac [35]
Additional file 10 <i>Globularia alypum</i> L. s.l. (PORUN - TTF2077) GLOBULARIACEAE	<i>zrega, zrga</i> (arab); <i>taselrha?</i> (berber); <i>tidi-n-tenet</i> (tuaregh)	Leaves branches Aerial part	The decoction is used as a laxative, against intermittent fevers and topically in the cure of furuncles	In North African countries, branch and leaf decoctions are prepared to cure intermittent fevers, arthritis and rheumatism and used as a depurative, diuretic, hypoglycemic and laxative [23,25]. The decoction of the aerial parts is administered against constipation, fever, and mycosis [28]. A leaf infusion is used as a hypoglycemic, digestive, and laxative as well as for bilious stimulation [24]
<i>Haplophyllum tuberculatum</i> (Forssk.) A. Juss. (PORUN - TTD59; TTF2064) RUTACEAE	<i>sezeret er rih</i> (arab)	Aerial part	Laxative	In Egypt and Saharan territories, flowering and fruiting branches are used to cure gastrointestinal affections, intermittent fevers, and rheumatism. The plant is also an aphrodisiac and administered against eye and ear affections [23]
Additional file 11 <i>Hedynois cretica</i> (L.) Dum.Cours. (PORUN - TTF2312) ASTERACEAE	<i>sezeret er rih</i> (arab)	Aerial part	Against meteorism and to cure haon (a kind of influence)	Medicinal uses not described for Maghreb and neighboring countries. Known as an edible plant in different Mediterranean countries [36,37]
<i>Hyoscyamus muticus</i> L. subsp. <i>falezlez</i> (Coss.) Maire (PORUN - TTF2315) SOLANACEAE	<i>gungot</i> (arab); <i>falezlez</i> (arab); <i>afahiehle</i> (temahac); <i>bathim</i> , <i>buthima</i>	Aerial part	The plant is severely poisonous. The extract of the aerial part mixed with butter is used externally against rheumatic affections	Known as medicinal plant in Marmarica [22]. In North Sahara, a potion is drunk as a tonic [29]. The oil macerate of leaves is topically used in Tassili N'ajjer against backache, muscular cramps, spasms, and palpitation anxiety as well as to treat eye inflammation and lice [28]
<i>Jateorhiza palmata</i> (Lam.) Miers (PORUN - TTD69; TTF2322) MENISPERMACEAE	<i>zaramba</i>	Roots	Infusion in orange water is given against cardiac affections	Its use is largely diffused in East and Central Africa as a bitter tonic, analgesic, and against diarrhea [38]
Additional file 12 <i>Juniperus oxycedrus</i> L. s.l. (PORUN - TTF2327) CUPRESSACEAE	<i>taga</i> (Algeria)	Wood of stem	Trunks are carbonized to give an oil (the cade oil) that is used against skin affections	In Tunisia and Morocco, the tar produced by the wood is antiparasitic and antiseptic for the skin [23]
<i>Launaea quercifolia</i> (Desf.) Pamp. (PORUN - TTF654) ASTERACEAE	<i>machinan</i> (arab)	Aerial part	Eaten against rheumatic aches	Medicinal uses not described for Maghreb and neighboring countries. In Morocco, <i>L. arborescens</i> is used to cure diabetes and against nausea and skin troubles [21]

Laurus nobilis L. (PORUN - TTD47)
LAURACEAE

Additional file 13

Dried leaves

Medicinal plant (uses not described). Dried leaves are sold in the markets by herbalists

Leaf and fruit oil has a cosmetic application for face care. An infusion of leaves and fruits is administered for dental hygiene and to cure liver, pancreas and digestive diseases. The decoction of the same parts is given to treat rheumatic pains [24,25]

Lawsonia inermis L. (PORUN - TTF1798)
LYTHRACEAE

hénna (arab); *alen* (berber); Aerial part
anela (temahac)

Astringent, vulnerary, used against dandruff and chilblains. Abortive

Leaves are antiseptic and astringent as well as used against eye affections and in the preparation of antirheumatic liniments. Flowers have insecticide properties [23]. Leaves powdered and mixed with water are applied as a hair tonic or mixed with lime juice as an emetic [24]. Also administered against gastric ulcer and kidney stones [25]

Lepidium sativum L. s.l. (PORUN - TTD34; TTF852)
BRASSICACEAE

habb' rsciad (arab); *carabau*, Seeds
tsc'uit (Algeria)

Sold in the markets and used to cure cough and asthma

In North Africa, the seeds, crushed with honey, or seed flour are administered against cough and pulmonary affections, delivery difficulties, heart tonic, revulsive and also in the cure of skin troubles. The leaves are considered tonic and effective in the prevention of scurvy; frequently used as a condiment [23,25]. The milk infusion of seeds is used to cure migraines. Seeds boiled in oil are used to treat diarrhea. Powdered seeds are externally applied for skin ulcers and warts. Seeds are also part of a preparation used in the treatment of sexual impotence [24]

Additional file 14

Marrubium alysson L. (PORUN - TTF1423)
Marrubium vulgare L. (PORUN - TTF1429)
LAMIACEAE

Aerial part

The powdered plant, mixed with oil, is used for rheumatism

Flowers have insecticide properties [23]. A leaf and stem decoction is used to treat intestinal pains, cough, and colds. Crushed leaves are used against ear pains [26]. An infusion of the aerial parts is used as an antipyretic, expectorant, antidiarrheal, tonic, bilious stimulation and for bronchitis and menstrual pains. The aerial part is also boiled with wine to obtain a syrup used as a stomachic [24,25]

Matricaria aurea (Loefl.) Sch. Bip. (PORUN - TTD25; TTF666)
ASTERACEAE

Flowering branches

The decoction is used against gastro-intestinal affections

In the Middle East, the plant is used as a substitute for *M. chamomilla* [39]

Additional file 15

<i>Myrtus communis</i> L. s.l. (PORUN - TTD31) MYRTACEAE Additional file 16	<i>rehân, ghemmâm, gidra</i> (arab)	Leaves flowers	and uses	The plant has many medicinal uses	The plant is sold in the Algiers and Rabat markets for use against diarrhea, gastrointestinal disorders, asthma and other respiratory ailments. It is also topically applied for painful organs. The plant is also used in perfumery and cosmetics [23,25]. Leaves and buds are used against hemorrhoids and skin affections. Fruits are administered for the cure of ocular disorders. An infusion of the leaves is used as a mouthwash against gingivitis and in association with walnut and mulberry in the treatment of diabetes [21]. A leaf decoction is used against cardiac and intestinal affections [26]. A leaf infusion is used against pneumonia, diarrhea and to promote wound healing. For hair care, a mixture with <i>Lawsonia inermis</i> is locally applied [24]
<i>Nitraria retusa</i> (Forsk.) Asch. (PORUN - TTF2317) ZYGOPHYLLACEAE	<i>gârdagh</i> (arab); (temahac)	<i>atazzim</i> Leaves		Crushed leaves are put in hot water and then used as a poultice against swellings	
<i>Olea europaea</i> L. s.l. (PORUN - TTF1811) OLEACEAE	<i>zêt, zêt ez zitûna</i>	Oil		Excipient of many plant drugs	A leaf decoction is used against nervous troubles and as an anthelmintic. Powdered leaves are used in the treatment of diabetes; burnt leaves are used to cure eye affections. Olive oil is administered to cure dry cough and grippe [26]. In Morocco, it is used for mouth hygiene, stomach pains, intestinal diseases, and diabetes [24,25]
<i>Opuntia ficus-indica</i> (L.) Mill. (PORUN - TTF2329) CACTACEAE	<i>hîndî</i> (arab)	Flowers		Medicinal properties (not described)	The powdered flower is used against stomach disorders [26]. An infusion of flowers is administered to stop diarrhea and as a diuretic. Flowers are also part of a preparation used as a calefacient [24]. In Morocco, it is also administered to cure bladder, kidney and uterus infections [25]
<i>Orchis mascula</i> (L.) L. s.l. (PORUN - TTF2326) ORCHIDACEAE	<i>sahalep</i>	Whole plant		Used in the preparation of a medicinal powder	No medicinal use reported for Maghreb or Sahara
<i>Origanum majorana</i> L. (PORUN - TTD41) LAMIACEAE Additional file 17	<i>marđúscia</i> (arab)	Aerial part		Against dysentery	In the Algiers market, the plant is sold to cure eye affections [23] In Morocco, the infusion of branches is used to treat chills, fever, cough, and flatulence [25,26]
<i>Origanum vulgare</i> L. s.l. (PORUN - TTF2303) LAMIACEAE		Flowers		Flowers have medicinal properties	In Tunisia and Algeria, the leaves and flowering branches are used as a stimulant [23]
<i>Paliurus spina-christi</i> Mill. (PORUN - TTF2321) RHAMNACEAE	<i>sédr, zegregh, ennab, corna</i> (arab); <i>abaga, labacat</i> (temahac); <i>magaria, cussulu</i> (Algeria)	Young shoots Leaves		Anthelmintic	Medicinal uses not reported in Maghreb and neighboring regions. Used in the East Mediterranean region to treat respiratory, circulatory and gastro-intestinal disorders [40]

<i>Paronychia argentea</i> Lam. (PORUN - TTF2301) PARONYCHIACEAE	<i>theia el arab, theia el gebel</i> (Algeria)	Aerial part	Infusion used against dysentery	Used as an antidiabetic in the East Mediterranean [41]
<i>Peganum harmala</i> L. (PORUN - TTD35; TTF2067) ZYGOPHYLLACEAE Additional file 18	<i>hármal</i> (arab); <i>bender tifi, bender tiffin</i> (tuaregh)	Seeds	Oil is used against headaches; burnt against mental diseases	The plant is sold in North African markets and used for a large number of affections. In Sudan, it is used against inflammations of the head, face and mouth as well as against headache and sinusitis [31]. The leaf decoction is used against high blood pressure and hemorrhoids [21]. In Tassili N'ajjer (Algeria), a decoction of seeds is taken to treat a large number of affections, ranging from skin diseases to nervous disorders, including anxiety. Also used to treat diabetes, heminthiasis and jaundice. The seed decoction is externally applied against tumors, eczema and lice [28]. Similar uses are also reported for Morocco and the Northern Sahara [25,29]. Powdered seeds of barley and <i>P. harmala</i> are topically applied against rheumatism [21]
<i>Periploca angustifolia</i> Labill. (PORUN - TTF2306) APOCYNACEAE	<i>sinuâc, teborac, arac</i> (arab); <i>têhac</i> (temahac); <i>chigu</i> (near Tchad)	Dried leaves	A decoction of dried leaves is used against syphilis mixed with a substance called ras el hânut	The seed decoction is used as a local analgesic and antirheumatic [23]. In Tassili N'ajjer (Algeria), the decoction of seeds or the aerial part of the plant can be drunk as an abortive and to cure diabetes. Externally, used to treat rheumatism and various pains [28]. Fodder appreciated in arid areas, eaten by camels, sheep and goats
<i>Phoenix dactylifera</i> L. (PORUN - TTF2323) ARECACEAE	<i>na'ha</i> (arab); <i>tezdit</i> (berber); <i>tazzeit, tazeit</i> (temahac); <i>zui</i> (Augila)	Fruits	Fruits	In North African countries, the wood of the palm is used as toothbrush. Dates are used as against ulcers of the genital organs and as a diuretic, laxative, and tonic [23]
<i>Piper retrofractum</i> Vahl (PORUN - TTD19) PIPERACEAE Additional file 19	<i>dahr el filfil</i>	Flowers	Imported in Libya from Sudan. When mixed with honey is used as an aphrodisiac	The plant is imported from Asian countries. In Morocco, the plant is known for its aphrodisiac, caleficient and magic properties [25]
<i>Pistacia atlantica</i> Desf. (PORUN - TTD3; TTF16) ANACARDIACEAE Additional file 20	<i>batûm</i> (arab); <i>tizert</i> (berber)	Fruits	Sold in the markets. It is chewed against respiratory affections. The oil has similar properties	Leaves are used against skin affections. Fruits are used to season dates. Used for tanning [23]. In Marmarica, it is used as fuel, grazing and medicinal plants [22]. Burnt leaves are used in a poultice against eye affections [26]. In Morocco, fruits are administered against stomach-ache, whereas the gall is used for cosmetic applications, against fever and stomach diseases [25]

<i>Plantago afra</i> L. s.l. (PORUN - TTF2324) PLANTAGINACEAE	<i>anâm, nenâm</i> (arab)	Whole plant	Vulnery. Powdered and dried is administered in topical applications	In Marmarica, used as medicinal plant [22]. In North African countries, used against metabolic disorders, gastro-intestinal affections, hemorrhoids, skin diseases, urinary tract disorders, and venereal diseases. Fresh leaves are applied topically for poison ivy, insect bites and stings [21]
<i>Punica granatum</i> L. (PORUN - TTD37) PUNICACEAE Additional file 21	<i>rummân, rummâna</i> (arab); <i>taarmunt, armun</i> (fruit) (berber); <i>tarrumant</i> (temahac)	Flowers	Flowers are medicinal. Seeds are tonic and aphrodisiac	In Egypt, fruit peels are sold as an astringent; dried powdered peel is sold in Morocco as an antidiabetic, antidiarrheal, antiseptic, antitumor, astringent, and hemostatic as well as against gastro-intestinal and gynecological disorders and for cleansing the teeth [23,24,25,26]
<i>Ricinus communis</i> L. s.l. (PORUN - TTD14) RUTACEAE Additional file 22	<i>chêrua</i> (arab)	Seeds	The water infusion is a laxative (also chewed seeds). Leaves are used in the preparation of poultices	Known as a medicinal plant in Marmarica [22]. In Central Sahara, the seed decoction is used against fevers and headaches or externally applied to cure trachoma, aphthae, and hair loss [28]. A decoction is administered to treat cow jaundice [24]. In Sudan, the fresh leaves are rubbed on the head to relieve headache or on the legs against swellings [30]
<i>Rosa damascena</i> Mill. (PORUN - TTF2325) ROSACEAE	<i>uârd</i> (arab)	Rose buds	The herbalists in Tripoli sell dried rose buds for medicinal purposes (sciûs el uârd)	Dried flower buds are used against headache, stomach pains, toothaches, and as laxative and hair tonic are sold in North African markets [23,25]. Used also in the cure of numerous affection of the eyes and ears [23] The flower infusion is used as a laxative. A mixture with other plants is applied externally [24]
<i>Rosmarinus officinalis</i> L. (PORUN - TTD16; TTF1441) LAMIACEAE Additional file 23	<i>clîl</i> (arab); (berber)	Leaves	The infusion is used against cough; ground dried leaves in oil are vulnerary	The leaf decoction is administered against intestinal parasites and rheumatism in Central Sahara [26] as well as an emmenagogue, spasmolytic against gastro-intestinal and liver disorders, diuretic, carminative and sedative in Morocco [25]. A leaf infusion is also prepared against tachycardia and as a cholagogue and vasopressor. Leaves are also externally applied against wrinkles, muscular pains and rheumatism and as a vulnerary [24]. In Tunisia, rosemary leaves are used as an antispasmodic for the digestive tracts and as a vermifuge. Dried leaves, ground and mixed with olive oil, are put on recent circumcision wounds [21]

<i>Ruta chalepensis</i> L. (PORUN - TTF2069) <i>Ruta</i> sp. (PORUN - TTD24) Additional file 24	<i>fgél, figél</i> (arab); (temahac) <i>issín</i>	Aerial part	Sold in Tripoli market, and used against rheumatic affections and echymosis. The smell of the plant keeps the scorpions away from houses In Rabat Market, the plant is sold for use against nervous affections, and in Algiers markets, for use against vomiting and fevers of children and babies [23]. Washing with crushed leaves is used against ear pains, and leaves are smoked to keep away bad spirits [26]. In Morocco, the aerial parts are used as an abortive, for intestinal and hepatic diseases, male sterility, and vitiligo [25]. Also known as a vulnerary, emmenagogue, and spasmodic; the fresh plant is used as a scorpion and insect repellent. Leaves and seeds, boiled in olive oil, are rubbed on the skin to treat rheumatism and swellings [21]
<i>Ruta graveolens</i> L. (PORUN - TTF2309) RUTACEAE	<i>fgél, fesál</i> (arab)	Aerial part	In Cyrenaica, a potion made with this plant is administered to newborns as a tonic
<i>Salvia aegyptiaca</i> L. (PORUN - TTF1370) LAMIACEAE	<i>ra'al, sezeret el rházel</i> (arab, ex Muschler); <i>safsaf</i> (tuaregh)	1. aerial part; 2. leaves	The infusion is digestive. The leaves are kept in the nose to give a fresh sensation
<i>Schoenocaulon officinale</i> (Schltdl. et Cham.) A. Gray ex Benth. (PORUN - TTF2311) MELANTHIACEAE	<i>duá ghémel</i>	Seeds	Against lice
<i>Smyrniolum olusatrum</i> L. (PORUN - TTF2193) APIACEAE	<i>calch</i> (arab)	Aerial part	Abortive
<i>Tanacetum parthenium</i> (L.) Sch. Bip. (PORUN - TTF2332; TTD63) ASTERACEAE Additional file 25	<i>usciach</i>	Gum resin	Imported from Marseille and sold in the markets. It is dissolved in vinegar against contusions or to cure furuncles
<i>Teucrium polium</i> L. s.l. (PORUN - TTD65; TTF1389) LAMIACEAE Additional file 26 <i>Thapsia garganica</i> L. s.l. (PORUN - TTF2196) BRASSICACEAE	<i>giaád, zaád</i> (arab); <i>techmezzutin</i> (tuaregh) <i>drias</i> (arab)	Roots Aerial part Inflorescence	Medicinal properties (not described) Gastro-intestinal affections Used in North Africa against dysmenorrhea [21]. In Morocco, the aerial parts are used against chill, edema, live pain and for blood cleansing [25] The infusion of the aerial parts is used against cough and rheumatic pains. The root infusion is used to treat liver and bladder diseases, and an oil is externally applied for swelling and wrinkles [24]

Thymus capitatus (L.) Hoffmanns. et Link (PORUN - TTD70; TTF1401)
Additional file 27

Thymus mastigophorus Lacaita (PORUN - TTF2328)
 LAMIACEAE
Trigonella foenum-graecum L. (PORUN - TTF1596)
 FABACEAE

zaátar (arab); sótar (berber)	Leaves and floral shoots	A cold infusion against cough	Eaten as vegetable in Marmarica and also known as medicinal plant [22]. In Libya, the plant is used for coughs, as a tonic, and against skin affections [23]
zaátar (arab); arrar? (berber)			
héiba (arab)	Seeds	Against cough, febrifuge	Cultivated and naturalized in North Africa [23]. The seed decoction is used for uterine pains. Boiled seeds are also ingested as a hypoglycemic, while powdered seed mixed with water are externally applied as a hair tonic [24]. In Morocco, seeds are considered to be blood cleansing and an aortic-palpitation reconstituant [25]. In Sudan, it is administered against rheumatism and dysentery as well as for cleaning the blood. It is also reputed as a lactagogue [31]. In Tassili N'ajjar (Algeria), the decoction of the aerial parts or seeds are used to cure diabetes, clean the blood, and as a tonic and an analeptic [28]. Seed maceration is effective to treat diabetes, scurvy, and digestive troubles [26]. Used in North Africa for a wide range of affections [23]. In Sudan, used for colds and rheumatism as well as to treat pneumonia [31]
<i>Zingiber officinale</i> Roscoe (PORUN - TTF2330) ZINGIBERACEAE	Roots	Aphrodisiac	

Evernia furfuracea (L.) Mann (PORUN scíba - TTD56)
 PARMELIACEAE

Additional file 28
Ramalina calicaris (L.) Fr. (PORUN - TTD53)
 RAMALINACEAE
Additional file 29

Ramalina farinacea (L.) Ach. (PORUN scíba - TTD54)
 RAMALINACEAE

Additional file 29
Usnea plicata (L.) Fries (PORUN - TTD55)
 USNEACEAE
Additional file 30

The occurrence of ? after a local name of a plant is a Trotter indication.

Additional files

- Additional file 1.** *Ajuga iva* (L.) Schreb. (PORUN - TTD52), aerial part.
- Additional file 2.** *Alpinia officinarum* Hance (PORUN - TTD17), rhizome.
- Additional file 3.** *Artemisia arborescens* L. (PORUN - TTD51), young shoots, flowers and leaves.
- Additional file 4.** *Boswellia sacra* Flüeckiger (PORUN - TTD39), resin of stem.
- Additional file 5.** *Capparis orientalis* Veillard in Duh. (PORUN - TTD7), leaves.
- Additional file 6.** *Cuminum cyminum* L. (PORUN - TTD9), fruits.
- Additional file 7.** *Cupressus sempervirens* L. (PORUN - TTD60), cones and seeds.
- Additional file 8.** *Cymbopogon schoenanthus* (L.) Spreng. s.l. (PORUN - TTD79), basal part of the plant.
- Additional file 9.** *Elettaria cardamomum* (L.) Maton (PORUN - TTD8), fruits.
- Additional file 10.** *Ferula marmarica* Asch. et Schweinf. (PORUN - TTD22), gum resin.
- Additional file 11.** *Haplophyllum tuberculatum* (Forssk.) A. Juss. (PORUN - TTD59), aerial part.
- Additional file 12.** *Jateorhiza palmata* (Lam.) Miers (PORUN - TTD69), roots.
- Additional file 13.** *Laurus nobilis* L. (PORUN - TTD47), leaves.
- Additional file 14.** *Lepidium sativum* L. s.l. (PORUN - TTD34), seeds.
- Additional file 15.** *Matricaria aurea* (Loefl.) Sch. Bip. (PORUN - TTD25), flowering branches and leaves.
- Additional file 16.** *Myrtus communis* L. s.l. (PORUN - TTD31), leaves and flowers.
- Additional file 17.** *Origanum majorana* L. (PORUN - TTD41), aerial part.
- Additional file 18.** *Peganum harmala* L. (PORUN - TTD35), seeds.
- Additional file 19.** *Piper retrofractum* Vahl (PORUN - TTD19), flowers.
- Additional file 20.** *Pistacia atlantica* Desf. (PORUN - TTD3), fruits and gall-nut.
- Additional file 21.** *Punica granatum* L. (PORUN - TTD37), flowers.
- Additional file 22.** *Ricinus communis* L. s.l. (PORUN - TTD14), seeds.
- Additional file 23.** *Rosmarinus officinalis* L. (PORUN - TTD16), leaves.
- Additional file 24.** *Ruta* sp. (PORUN - TTD24), aerial part.
- Additional file 25.** *Tanacetum parthenium* (L.) Sch. Bip. (PORUN - TTD63), roots.
- Additional file 26.** *Teucrium polium* L. s.l. (PORUN - TTD65), aerial part.
- Additional file 27.** *Thymus capitatus* (L.) Hoffmanns. et Link (PORUN - TTD70), leaves and floral shoots.
- Additional file 28.** *Evernia furfuracea* (L.) Mann (PORUN - TTD56), thallus.
- Additional file 29.** *Ramalina calicaris* (L.) Fr. (PORUN - TTD53) and *Ramalina farinacea* (L.) Ach. (PORUN - TTD54), thallus.
- Additional file 30.** *Usnea plicata* (L.) Fries (PORUN - TTD55), thallus.




Figure 1



Golfo di Gabes (Piccola Sirte)

Golfo di Sidra (Gran Sirte)

BARCA
CIRENAICA

S
I
A

Tripoli
SABIA
TRIPOLI
P
O
L
I

Hammada el Homra
(Doveria Pietrosa Altipiana)

Dune di sabbia d'Eddeyen

F
E
Z
Z
A
N

Hammada di Murzuk

Murzuk

D
E
S
E
R
T
O
Serir
D
I
S
erir
L
I
B
I
A


David di Kufra

E
G
I
T
T
O

Segni Convenzionali

- Confine di Stato
- Grande Circonferenza (meridiano)
- Circolo di latitudine
- Città di oltre 100.000 abitanti
- Città di meno di 100.000 abitanti
- Linee di Riferimento Geografiche
- Semplici costruzioni
- Semplici costruzioni per segnalare luce a Torretta

N.B. L'isolezione dei paesi è secondo l'opinione dei compilatori.



10° 15° 20° Longitudine dal meridiano di Greenwich

MISSIONE AGRICOLA PER LO STUDIO DELLA TRIPOLITANIA

HERBARIUM PORTIDENSE

Argemone siliqua var. *argemone* L. (Kunt.) Voss
1/2 m. 10. 1. 1900. 2. 1900.

MISSIONE AGRICOLA PER LO STUDIO DELLA TRIPOLITANIA

HERBARIUM PORTIDENSE

Argemone siliqua L. (Kunt.) Voss
1/2 m. 10. 1. 1900. 2. 1900.

Figure 4



Figure 5



Figure 6

HERBARIUM PORTUENSE



105, Chól

Antimonas - cobalt.

o. Algenberg



Figure 7

HERBARIUM PORTIGENSE



HERBARIUM PORTIGENSE



114. "Tin"



HERBARIUM PORTIGENSE

ad Engila a sud di Rio Zava

argilla smaltina per lavare impregnata
anche in biondo come impregnata per
le parti in...

HERBARIUM PORTIGENSE

Figure 8

Additional files provided with this submission:

Additional file 1: Additional file 1.tif, 327K
<http://www.ethnobiomed.com/imedia/1314604212663236/supp1.tif>

Additional file 2: Additional file 2.tif, 288K
<http://www.ethnobiomed.com/imedia/3993809966632377/supp2.tif>

Additional file 3: Additional file 3.tif, 313K
<http://www.ethnobiomed.com/imedia/1616786922663237/supp3.tif>

Additional file 4: Additional file 4.tif, 313K
<http://www.ethnobiomed.com/imedia/1380677171663237/supp4.tif>

Additional file 5: Additional file 5.tif, 341K
<http://www.ethnobiomed.com/imedia/2126820826663238/supp5.tif>

Additional file 6: Additional file 6.tif, 350K
<http://www.ethnobiomed.com/imedia/8291739436632385/supp6.tif>

Additional file 7: Additional file 7.tif, 292K
<http://www.ethnobiomed.com/imedia/1110587725663238/supp7.tif>

Additional file 8: Additional file 8.tif, 325K
<http://www.ethnobiomed.com/imedia/1922301846632395/supp8.tif>

Additional file 9: Additional file 9.tif, 216K
<http://www.ethnobiomed.com/imedia/6633236206632387/supp9.tif>

Additional file 10: Additional file 10.tif, 256K
<http://www.ethnobiomed.com/imedia/5259694096632387/supp10.tif>

Additional file 11: Additional file 11.tif, 322K
<http://www.ethnobiomed.com/imedia/1329405231663239/supp11.tif>

Additional file 12: Additional file 12.tif, 256K
<http://www.ethnobiomed.com/imedia/5741473676632398/supp12.tif>

Additional file 13: Additional file 13.tif, 288K
<http://www.ethnobiomed.com/imedia/1755453590663238/supp13.tif>

Additional file 14: Additional file 14.tif, 307K
<http://www.ethnobiomed.com/imedia/1148267851663239/supp14.tif>

Additional file 15: Additional file 15.tif, 338K
<http://www.ethnobiomed.com/imedia/7084850546632389/supp15.tif>

Additional file 16: Additional file 16.tif, 392K
<http://www.ethnobiomed.com/imedia/1388386553663239/supp16.tif>

Additional file 17: Additional file 17.tif, 348K
<http://www.ethnobiomed.com/imedia/1355496916663239/supp17.tif>

Additional file 18: Additional file 18.tif, 148K
<http://www.ethnobiomed.com/imedia/4063821486632390/supp18.tif>

Additional file 19: Additional file 19.tif, 235K
<http://www.ethnobiomed.com/imedia/2130013224663239/supp19.tif>

Additional file 20: Additional file 20.tif, 340K
<http://www.ethnobiomed.com/imedia/6430357276632391/supp20.tif>

Additional file 21: Additional file 21.tif, 310K
<http://www.ethnobiomed.com/imedia/1681840858663239/supp21.tif>

Additional file 22: Additional file 22.tif, 218K
<http://www.ethnobiomed.com/imedia/7494888966324013/supp22.tif>

Additional file 23: Additional file 23.tif, 251K

<http://www.ethnobiomed.com/imedia/2640019136632392/supp23.tiff>

Additional file 24: Additional file 24.tif, 234K

<http://www.ethnobiomed.com/imedia/8502910546632401/supp24.tiff>

Additional file 25: Additional file 25.tif, 188K

<http://www.ethnobiomed.com/imedia/1168493828663240/supp25.tiff>

Additional file 26: Additional file 26.tif, 357K

<http://www.ethnobiomed.com/imedia/1655101572663239/supp26.tiff>

Additional file 27: Additional file 27.tif, 284K

<http://www.ethnobiomed.com/imedia/1113360926632402/supp27.tiff>

Additional file 28: Additional file 28.tif, 363K

<http://www.ethnobiomed.com/imedia/9128639466323933/supp28.tiff>

Additional file 29: Additional file 29.tif, 331K

<http://www.ethnobiomed.com/imedia/1411018733663240/supp29.tiff>

Additional file 30: Additional file 30.tif, 241K

<http://www.ethnobiomed.com/imedia/4986425536632393/supp30.tiff>