



Ethnoknowledge of Bukusu community on livestock tick prevention and control in Bungoma district, western Kenya

Wycliffe Wanzala^{a,b,*}, Willem Takken^c, Wolfgang R. Mukabana^d, Achola O. Pala^e, Ahmed Hassanali^{b,f}

^a School of Pure and Applied Sciences, Department of Biological Sciences, South Eastern University College (A Constituent College of the University of Nairobi),

P.O. Box 170-90200, Kitui, Kenya

^b Behavioural and Chemical Ecology Department (BCED), International Centre of Insect Physiology and Ecology (ICIPE), P.O. Box 30772-00100 GPO, Kasarani, Nairobi, Kenya

^c Wageningen University and Research Centre, Laboratory of Entomology, P.O. Box 8031, 6700 EH Wageningen, The Netherlands

^d School of Biological Sciences, University of Nairobi, P.O. Box 30197-00100, Nairobi, Kenya

^e Technology Transfer Unit, International Centre of Insect Physiology and Ecology (ICIPE), P.O. Box, 30772-00100 GPO, Kasarani, Nairobi, Kenya

^f School of Pure and Applied Sciences, Kenyatta University, P.O. Box 43844-00100 GPO, Nairobi, Kenya

ARTICLE INFO

Article history:

Received 24 August 2011

Received in revised form

14 December 2011

Accepted 13 January 2012

Available online 24 January 2012

Keywords:

Bukusu

Ethnoveterinary knowledge

Mixed pastoral/arable farmers

Tick ethnocontrol practices/methods

Anti-tick ethnobotanicals

ABSTRACT

Ethnopharmacological relevance: To date, nomadic communities in Africa have been the primary focus of ethnoveterinary research. The Bukusu of western Kenya have an interesting history, with nomadic lifestyle in the past before settling down to either arable or mixed arable/pastoral farming systems. Their collective and accumulative ethnoveterinary knowledge is likely to be just as rich and worth documenting.

Aim of the study: The aim of the present study was to document indigenous knowledge of the Bukusu on the effect of livestock ticks and ethnopractices associated with their management. It was envisaged that this would provide a basis for further research on the efficacy of these practices that could also lead to the discovery of useful tick-control agents.

Materials and methods: Non-alienating, dialogic, participatory action research (PAR) and participatory rural appraisal (PRA) approaches involving 272 women and men aged between 18 and 118 years from the Bukusu community were used.

Results: Ticks are traditionally classified and identified by colour, size, host range, on-host feeding sites, and habitat preference. Tick-associated problems recognised include *kamabumba* (local reference to East Coast fever, Anaplasmosis or Heartwater diseases transmitted by different species of livestock ticks) and general poor performance of livestock. Traditional methods of controlling ticks include handpicking, on-host use of ethnobotanical suspensions (prepared from one or more of over 150 documented plants) to kill the ticks and prevent re-infestation, fumigation of infested cattle with smoke derived from burning ethnobotanical products, burning pastures, rotational grazing ethnopractices, and livestock quarantine. **Conclusions:** The study confirms that the Bukusu have preserved rich ethnoveterinary knowledge and practices. It provides some groundwork for elucidating the efficacy of some of these ethnopractices in protecting livestock from tick disease vectors, particularly those involving the use of ethnobotanicals, which may lead to the discovery of useful ant-tick agents.

© 2012 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

Bungoma district is located in western province of Kenya between latitude 00°28' and 10°30'N and longitude 34°20' and 35°15'E (Fig. 1). It occupies an area of 3074 km² on the southern slopes of Mt. Elgon (Fig. 1), with a population of about 2 million

* Corresponding author at: School of Pure and Applied Sciences, Department of Biological Sciences, South Eastern University College (A Constituent College of the University of Nairobi), P.O. Box 170-90200, Kitui, Kenya. Tel.: +254 02 2413859; mobile: +254 724 396358/733 258331; fax: +254 02 4445763.

E-mail address: osundwa1@yahoo.com (W. Wanzala).

people. It lies in an agro-ecological zone stretching from Tropical Alpine Zone to Lower Midland Zone with tea, wheat/maize, pyrethrum, coffee, sunflower, maize, sugarcane and cotton plantations (Martina, 1998, 2001). The predominant off-farm vegetation patterns are riverine forests, rocky forested hillsides, hedgerows, wooded grassland relicts, woodlands or colline forest relicts and tree groves whereas the noticeably tree rich on-farm management units are home gardens, homesteads, live fences, coffee and banana groves and annual cropping fields (Martina, 2001).

The Bukusu community (also known as *Babukusu* and/or *Bakitoshi*) represents one of 19 ethnic groups (sub-tribes) of the Luhya people of interlacustrine Bantu group of East Africa (Central Bureau




Fig. 1. The location of Bungoma district in Kenya. Note the positioning of the district along the southern slopes of Mt. Elgon.

of Statistics, 1996) and are dominant in the Bungoma district (Fig. 1). Although arable farming is now the main occupation of the community, livestock farming is still held in high esteem and livestock products are highly valued in all traditional and cultural practices. To date, research on ethnoveterinary knowledge has focused mainly on pastoral and/or nomadic communities (ITDG and IIRR, 1996). Settled or mixed farming communities such as the Bukusu have been neglected and their ethnoveterinary practices are at risk of being lost (Kofi-Tsekpo and Kioy, 1998). Almost all of these communities, the Bukusu included, largely led nomadic lifestyle before settling down to either arable or mixed arable/pastoral farming systems (Makila, 1978). Their collective and accumulative ethnoveterinary knowledge is likely to be just as rich as that of nomadic communities (ITDG and IIRR, 1996).

In this study, we discuss the results of a survey conducted to document and analyse anti-tick ethnoveterinary practices of the Bukusu community in western Kenya. The study demonstrates an important link between plant biodiversity and livestock healthcare ethnoknowledge systems in the Bukusu community.

1.1. Bungoma district and the Bukusu community

Bungoma is a district with a wide range of ethnic diversity of populations in western Kenya (Fig. 1). This is partly due to its establishment as the last stop on the Kenya–Uganda railway in the 1920s, rapidly growing into a complex commercial and colonial and post-colonial administrative centre along the railway line

(Port, 2000). However, the majority of the residents of Bungoma still belong to the indigenous Bukusu, who have lived near the border with Uganda along the southern slopes of Mt. Elgon ever since their migration from the Sudan/Egypt border during the second half of the last century (Were, 1967; Makila, 1978). The Bukusu community comprises 16.35% of the Luhya population (Central Bureau of Statistics, 1996). Their cultural and traditional life has been heavily shaped in time and space by immigrants of both Bantu and Nilotic origin (Makila, 1978; Wandibba, 1998; Roach, 2003). Members of the Bukusu community hold livestock keeping and livestock per se in high esteem, as livestock is used as a measure of one's wealth, power, authority and societal class status (Kenya National Archives, 1954; Nasimiyu, 1985; Nangendo, 1994). Moreover, the animals' blood and meat are highly valued in different traditional and cultural practices, including funeral ceremonies (where cattle skin is used as a coffin), circumcision, agricultural festivals, bride wealth exchanges, and during cleansing ceremonies and sacrifices to appease the spirits (Were, 1967; Makila, 1978). Plants have been highly valued for their nutritional and economic value, as well as in ethnomedicinal and veterinary applications, traditional education and socio-cultural life (Martina, 1998; Mann, 2001; UNEP World Conservation Monitoring Centre, 2003). For centuries, plants have been part of the Bukusu's land-use system and agro-ecological layout. This is reflected in the practice of selective clearing of trees, forests, and bushes, leaving trees and woodlots that are perceived to be useful to grow in the arable fields for fodder, timber, food, medicines, windbreaks,

fences/boundaries, supporting crops, shrines, etc. (Martina, 1998; Mann, 2001). A significant proportion of Bukusu poems, riddles, and proverbs touch on the symbolic value attached to indigenous plants, in daily life experiences as well as in philosophical reflections, which elders considered to be essential skills to equip growing children (Makila, 1978; Wandibba, 1998; Mweseli, 2004). Some plants were not put to regular use, but were specifically preserved for ancestors or for certain functions and cultural occasions (Martina, 1998). A great deal of information regarding the use, management, and socio-cultural roles of plants has been preserved by the older generation and this has been considered a useful basis of community-specific conservation strategy (Juma, 1989).

Although the Bukusu continue to maintain a strong attachment to their culture, traditions and superstitions (Corbitt and Wanyama, 2004), there is evidence of gradual erosion of their ethnoscientific knowledge base and ethnopractices (Wandibba, 1998; Roach, 2003; Corbitt and Wanyama, 2004). This has risen from a combination of factors. First, as sedentary pastoralists, they have had more time for arable farming at the expense of their original livestock farming. Early in the last century, they lived mainly on millet, cassava and livestock, practising mixed farming with arable farming bias. Now they have diversified to maize, beans, potatoes, bananas and cabbage for food, and sugarcane, cotton, tobacco and sunflower for cash. This expanded range of arable farming activities is currently dominating land use and threatens the survival of grasslands, woodlands, bushes and forests, the source of ethnopharmacologically active agents upon which ethnoknowledge is based (Biosafety News, 2002). Other factors contributing to the erosion of ethnoknowledge of the Bukusu include: (1) untimely deaths of persons with ethnoknowledge without this being documented, (2) extinction of specific plant and animal species and ritual practices, (3) encroachment of development and modernization on cultural and traditional life, (4) adoption of lifestyles and education systems that do not embrace ethnoknowledge, (5) shifting bias in religious beliefs, (6) perception of certain socio-cultural practices as unhygienic and satanic, and (7) cost- and health-related risks involved in certain socio-cultural practices. The result is that either little or no information is being passed on to the next generation (Wandibba, 1998; Roach, 2003; Corbitt and Wanyama, 2004).

1.2. The significance of archiving and valuing ethnoveterinary knowledge

The pastoral lifestyle was initially the mainstay of the Bukusu economy before arable farming overtook it, as evidenced by the rich archaeology, nomadic and pastoral vocabulary, traditions and socio-cultural anthropology of the Bukusu people (Makila, 1978). The community has a long history of ethnoveterinary practices in curing, preventing, controlling and managing livestock diseases and vector-borne problems. A rich set of anti-tick ethnoveterinary practices has evolved to avert tick-related losses and promote livestock survival in a tick-prone environment (Lewis, 2001). These ethnopractices have withstood the test of time and have a number of advantages over the modern equivalents. They integrate different health-based tactics that are cost-effective, easily applied, locally suitable, easily accessed, and environmentally friendly (Mathias-Mundy and McCorkle, 1989; Hamburger et al., 1991; Martin et al., 2001). Moreover, these community-specific ethnopractices are an integral part of the people's daily lives and are based on experience gained over centuries and adapted to local culture and environment. Ethnoknowledge is largely transmitted orally through demonstrations, songs, poems, drawings, paintings and stories from generation to generation (Kokwaro, 1993; Mweseli, 2004), and is only sketchily recorded in books (Abegaz and Demissew, 1998). In certain communities, the collective knowledge is believed to be owned by the ancestors and kept

under the custody of presently living elderly men and women whose mortality can lead to loss of potentially useful information.

This study was undertaken to survey, generate and document a database of anti-tick ethnopractices and knowledge amongst the Bukusu community. This resource may provide useful information for further scientific research, which can clarify the relative efficacies of different products and ethnopractices in controlling ticks and diseases they transmit (Alghali, 1992). In so doing, value-added knowledge can be provided back to the community in the form of useful products and services, and help ethnopractitioners gain incremental confidence in their own ethnoknowledge. This may help establish active links with the modern science network for sustainable use and management of local environment and its resources for improved animal productivity and livelihood.

2. Materials and methods

Before the start of this project, official permission was sought from the Government of Kenya (GOK) through the ministries of Education, Internal Security, Livestock Development, and the Department of Culture and Social Services. Through the ministry Education (Bungoma District Education Office) permission was sought to use upper primary school pupils (from standard 6 to 8) to identify and access some of the key respondents, particularly, ethnopractitioners (para-veterinarians). Prior informed consent was also sought from individual key respondents through the local administration in the office of the president, GOK.

2.1. The source of anti-tick ethnoknowledge

Anti-tick ethnoknowledge was surveyed from 13 different sources in the study area. The survey involved a sample of 272 key respondents of mixed sex and age (18–118 years old) who were identified through a number of sources, including the upper primary school questionnaire method. The ages of key respondents were confirmed from their: (1), birth certificates and (2), national Identity Cards (ID)/passports in Kenya. The sources included primary school pupils, who provided leads to key respondents, and local school teachers/education officers who helped select candidate schools, gave permission to distribute and explain the questionnaire to pupils, and helped to clarify the information being sought where possible. Local veterinarians, para-veterinarians, and agricultural extension officers responsible for providing extension services to farmers were also able to identify some key respondents. We also attended public meetings organized by local administrators to identify potential respondents. Local livestock traders and dealers, as well as individual livestock farmers, contributed their knowledge on tick control based on their professional and economic activities, whereas community/village/clan leaders/elders had information on communal tick control and management programmes funded by the Government and NGOs. Local ethnopractitioners, including traditional healers/herbalists/spiritualists/ritualists, formed a particular subset of knowledgeable people from whom key respondents were drawn. Other sources of data included NGOs such as the Council for Human Ecology-Kenya, the German Development Service and the Bungoma Indigenous Trees Conservation Club, which had prior experience and close interaction with local livestock farmers and had earlier documented part of their ethnoveterinary knowledge (Martina, 1998, 2001; Mann, 2001; Buteyo, 2004). Centres known for preparation of a local brew (locally known as *Busaa*) were the most important meeting points of old men and women and formed important venues for the discussion sessions. Secondary data were obtained from the Bungoma District Veterinary Office (DVO) records on livestock tick prevention and control. All these

groups were consulted because each was associated with a specific aspect of ethnoknowledge relevant to the study.

2.2. Key steps to accessing anti-tick ethnoknowledge

2.2.1. Constitution of a sampling group

The first step was the generation of a purposive sample of key respondents from the 13 sources mentioned above. Key respondents were local experts or people in the study area with knowledge of a particular issue or technology of interest (in this case, anti-tick ethnoknowledge) (Etkin, 1993; Waters-Bayer and Bayer, 1994; McCorkle et al., 1997). They have a more extensive understanding of local social and veterinary-cultural systems than others in the community. A purposive sample referred to a particular subset of knowledgeable people in the area of traditional control and management of livestock ticks. Intensive and extensive collaboration and interaction with these key respondents was considered to be an effective research strategy (Oakley, 1981; Warry, 1992). A random sample would not have been appropriate for this type of socio-cultural set-up, as not everyone sampled randomly may have the required knowledge (Etkin, 1993; Martin, 1996).

2.2.2. Local primary school pupils: the questionnaire method

This was the main method used to identify key respondents and obtain local and anti-tick ethnoknowledge from largely illiterate people. The use of children to collect such information has had precedents in the field (Lans and Brown, 1998; Lans, 2001), but there is very little published literature on this method (Campbell, 1994).

From a list of local primary schools at the District Education Office (DEO) in Bungoma, 10 schools were selected for participation in the survey studies. The selection of schools was based on variables such as rurality vs. urbanization, ethnicity, gender, ongoing teaching activities and geographical spread of the schools within the study area. In selected schools, pupils were visited and the basics of participatory research (Baldwin and Cervinskis, 1993) were explained, and the importance of their contribution was stressed. Pupils were asked to interview parents, guardians, relatives, friends, neighbours, etc. about traditional practices of tick control and to fill the answers of the respondents in a structured questionnaire. The questionnaire consisted of 15 questions: (1) the location where questionnaire is administered, (2) identity of the person being interviewed, (3) type and number of livestock kept by the person interviewed, (4) what ticks mean to the livestock farmer, (5) the kind of tick-related problems experienced by the livestock farmers, (6) responses to tick-related problems, (7) association between ethnoremedies and identified tick-related problem, (8) tick ethnocontrol remedies based on ethnobotanicals, (9) how these ethnobotanicals are harvested, processed, and applied, (10) when do livestock farmers apply their ethnoremedies, (11) mode of application of these ethnoremedies, (12) monitoring of cases after application, (13) any observations made regarding side-effects of these ethnoremedies, (14) any collaboration sought during tick prevention, control and management, and (15) personal observation of tick infestation made by the interviewer in the candidate livestock herd of the interviewee.

From the questionnaires, key respondents were identified based on whether the essay of the primary pupils and/or their responses indicated that a respondent had potentially useful information on anti-tick ethnopractices. This method was considered very useful and robust because it reduced the following sources of bias: (1) modeling bias, which was the projection of the interviewer's views on to those studied, (2) strategic bias, which was the expectation of benefits by the subject, (3) familiar relationships between interviewer and interviewee (children and parents/neighbours) which would reduce resistance to questioning but could lead to rote

answers and outsider bias (Sutton and Orr, 1991). These three pre-conceived notions would therefore lead to poor selection of key respondents (Etkin, 1993; Waters-Bayer and Bayer, 1994).

2.2.2.1. Personal interviews with selected key respondents. The third step involved interviews/discussions with the selected respondents. These were guided exchanges, semi-structured by a mental checklist of relevant points to confirm the validity of the information in the questionnaires.

2.2.2.2. Collection of plant specimens. Following a personal interview with the selected key respondents, a field trip was made to identify and collect the listed plant specimens and/or ethnobotanical products. The specimens were harvested, prepared, packaged and stored according to the herbarium rules and regulations until transported to the University of Nairobi Herbarium, Kenya for botanical identification using voucher specimens and according to the Hutchinson system of plant taxonomy based on the plants' probable phylogeny. While in the herbarium, further non-experimental studies were also conducted. For each plant species collected from the field, a voucher specimen was prepared and deposited in the Herbarium of the University of Nairobi, Kenya.

2.2.3. Focus-group discussions

The fourth step involved holding joint focus-group discussions with all stakeholders. In the study area, 11 focus-group discussions were formed, each comprising between 15 and 20 stakeholders based on their geographical location, ethnicity, conventional profession, economic activities, age, interest and practices of ethnohealth. A focus-group discussion was an exploratory discussion designed to obtain perceptions on a specific theme from a target group in a non-threatening environment (Krueger, 1988; Etkin, 1993). This kind of group interaction produced data and insights that would have otherwise been less accessible (Morgan, 1988). The interaction between all stakeholders formed the collaborative and non-alienating, dialogic, participatory action research (PAR) and participatory rural appraisal (PRA) approaches utilized to build a consensus and verify that the information from other interviewees was accurately recorded (IIRR, 1994). The group interaction also minimized the objectification of the respondents as the only source of data (Oakley, 1981). One purpose of this form of collaborative research was to shift decision making based on theoretical knowledge to the community, rather than conceding this role to the conventionally trained expert (Warry, 1992).

2.2.4. Collection of secondary data

The fifth step involved the collection of secondary data on anti-tick plants and ethnopractices from the District Veterinary Office (DVO) in Bungoma. This was followed by an extensive literature search on the taxonomy of the plant specimens collected and their ethnobotanical applications from the internet, livestock research institutions, non-governmental organizations (NGOs), East Africa and University of Nairobi herbaria libraries and laboratories.

2.2.5. Enumerations of documented plants with effects on livestock ticks

An extensive list of plants used as sources of ethnobotanicals, including their scientific and vernacular names, growth habits, family names and other information about their usage was prepared (Appendix 1). The plants are arranged according to their family names in an alphabetical order. Family taxonomic ranks and/or units are more stable than lower taxonomic levels such as genera and species and facilitate identification of new species particularly during field surveys (Sahney and Benton, 2008; Sahney et al., 2010). Because of the ethnic diversity amongst communities living in the study area (Bungoma County), more than one vernacular name may

be used to refer to a particular plant species and/or any other related plant species within a given genus or family. Two or more different plant species were found to have the same vernacular name depending on their geographical locations, uses, and associated ethnic group(s). In [Appendix 1](#), the classification of the plant specimens and/or ethnobotanical products into the column of growth life forms and/or habits was based on the definition and description of [Yumoto et al. \(1994\)](#).

3. Results and discussion

3.1. Local primary school pupil – the school questionnaire method

From the questionnaires administered to more than 300 pupils in 10 selected schools ([Fig. 1](#)), about 250 well-filled questionnaires were collected for evaluation of the anti-tick ethnoknowledge and 200 persons in the questionnaires were selected to form the key respondent team. The school method was instrumental in allowing the participation of a cross-section of local residents in the research survey. This helped us to obtain much useful information and ensured that plants with more than one vernacular name were correctly identified and their uses accurately recorded. In addition, it allowed a considerable number of illiterate individuals, ethnopractitioners and other key knowledgeable respondents living in remote areas to be reached and to meet the survey targets ([Sutton and Orr, 1991](#)). Many of these people gave useful leading viewpoints as shown in [Tables 1–4](#). Nevertheless, this approach was also realized to provide a bridge between elders and school pupils in terms of transmission of ethnoknowledge between generations. This had another great contribution in this study, because school schedules do not add local and indigenous knowledge into their curricula and school pupils rarely appreciate this knowledge from their illiterate parents, guardians, relatives and neighbours. But when they do provide such useful information for the research and it is well appreciated, the pupils recognise how elders of all ages kept and continue to keep important knowledge.

3.2. Interviews with key respondents and focus-group discussions

Focus-group discussions were instrumental in enabling us to transcribe the local Bukusu language accurately, especially with regard to local plant names and their meanings. The series of discussion sessions helped to harmonise amongst interviewees the use of multiple names referring to one plant and vice versa. A consensus was thus built on the identity of plants used locally, their local names, processing methods and uses. This facilitated the selection of the anti-tick plants that merited further scientific investigation. The experience with the respondents highlighted need for conventionally trained scientists to: (1) foster communication with different respondents and take time to establish a strong, trusting relationship based on openness and cooperation, (2) demonstrate respect for the ownership, source and origin of ethnoknowledge and the needs and sensitivities of its holders, and (3) undertake to provide value-added knowledge back to the community in the form of useful products (such as reports) and to share equitably with the key holders any benefits arising from the use of their ethnoknowledge. Specifically, ethnopractitioners need to be made aware of legislation protecting their ethnoknowledge base through relevant state ministries and possible threats from biopiracy.

3.3. Responses to questionnaires and results from the discussion groups

Despite initial constraints in accessing anti-tick ethnoknowledge, there was sufficient goodwill amongst most participating stakeholders to ensure substantial success in the research process.

Table 1a
Tick description by colour, size and life cycle type in relationship to on-host feeding site and habitat preference as a percentage of the responses ($n=987$).^a

Tick description	Site preference of the described ticks											NMAS	% total
	On host animals												
	Ear	Head	Dewlap/neck/brisket	Forelegs	Belly	Back/body sides	Hind legs	Tail/anal region	Udder/scrotum	Habitat			
										Bush	Grass		
Colour													
Red	5.37	1.32	2.43	1.52	0.51	1.11	1.62	2.74	3.34	0.00	0.51	0.30	20.77
Brown	5.88	1.93	3.34	2.13	0.61	1.01	2.23	4.05	4.56	0.00	0.30	0.30	26.34
Blue	6.08	2.13	1.95	2.03	0.61	1.11	2.13	3.55	4.76	0.00	0.41	0.10	24.86
Black	0.61	0.10	0.20	0.10	0.00	0.00	0.10	0.41	0.41	0.00	0.00	0.10	2.03
Multicolour	1.11	0.00	0.20	0.30	0.00	0.20	0.41	0.10	0.10	0.00	0.10	0.10	2.62
Grey	0.10	0.00	0.00	0.10	0.00	0.00	0.10	0.00	0.00	0.00	0.00	0.10	0.40
Yellowish	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.10
Size													
Small	1.32	0.10	0.51	0.41	0.20	0.10	0.51	0.30	1.01	0.00	0.00	0.41	4.87
Big	1.52	0.10	0.30	0.41	0.30	0.10	0.41	0.30	1.22	0.00	0.00	0.30	4.96
Life cycle type													
1-Host tick	0.71	0.41	0.30	0.00	0.00	0.00	0.20	0.61	0.51	0.20	0.20	0.10	3.24
2-Host tick	1.52	0.51	0.51	0.20	0.20	0.00	0.30	1.11	0.71	0.20	0.20	0.10	5.56
3-Host tick	1.11	0.51	0.41	0.10	0.20	0.00	0.20	0.61	0.71	0.10	0.10	0.20	4.25
% total	25.43	7.11	10.15	7.30	2.63	3.63	8.21	13.78	17.33	0.50	1.82	2.11	100.00

NMAS, no mention of association with any site of either host or habitat.

^a More than one response was received from each of the 272 respondents.

Table 1bObservation records by interviewees on the individual animals of their herds ($n = 233$).

Categories of observations	No. of herds observed	Mean no. of animals per herd	% of interviewees who made an observation
Many ticks (>20 per animal)	19	11 ± 1	8
Few ticks (~5 to 20 per animal)	21	10 ± 2	9
Very few ticks (<5 per animal)	16	8 ± 1	7
No tick was observed on any animals in the candidate herd	37	7 ± 1	16
No observation was made on individual animals in the candidate herd	140	10 ± 1	60

From the survey studies, it became apparent that although for a long time (since the coming on of conventional methods) the role of ethnoveterinary medicine and its potential contribution to livestock health have been neglected (Kofi-Tsekpo and Kioy, 1998), a large proportion of Bukusu people still depend on it for the welfare of their livestock. This was evidenced by the complex composition of the purposive sample ($n = 272$) of key respondents, and their voluminous and high-quality responses (Tables 1a and 1b). Even more important was the commitment and determination of all the stakeholders to have their local anti-tick ethnoknowledge recognised, documented and evaluated (Table 2). They also demonstrated interest in the results of scientific research on their ethnoknowledge and in any improvements that may be developed.

From discussions during one of the focus-group sessions held in Bungoma district under the auspices of Bungoma Indigenous Trees Conservation Club (BITCC) and Bungoma Herbalists Association (BHA), it was evident there was concern that the community could lose its rich socio-cultural heritage evidenced by lack of generalized use of this ethnobotanical knowledge across the Bukusu community (Wandibba, 1998; Corbitt and Wanyama, 2004; Buteyo, 2004). There was thus a consensus that efforts to conserve ethnobotanical knowledge of the Bukusu be intensified. Throughout the colonial era, indigenous knowledge was equated with witchcraft and wholly dismissed and actively suppressed. However, the use of this knowledge base continued in a clandestine manner to avoid punishment by the colonial administration, and only after independence did trends towards strengthening national and socio-cultural identities began to re-emerge (Mubukusu, pers. comm.). Succeeding governments encouraged integration of indigenous knowledge into existing health services. However, there was no clear legal framework for its operation. The conventionally trained doctors and various religious faith groups often took the advantage of lack of legal recognition of indigenous knowledge to denounce it. This attitude facilitated exploitation of ethnobotanical information without due recognition of indigenous source. Indeed, during our study, Bungoma ethnopractitioners complained about malpractices and biopiracy by researchers involved in bioprospecting, some of whom

reported their findings without giving the community appropriate recognition. This behavior has caused fear amongst ethnopractitioners and has contributed to their reluctance to share indigenous knowledge with foreigners. It also accounts for a discrepancy between knowledge availability and its use, as well as mistrust amongst stakeholders (Simon, 1998; Hempel, 2002). It is hoped that the present study, based on participatory research, will make some contribution towards reinforcing ownership and recognition of indigenous ethnobotanical knowledge amongst the Bukusu. Moreover, local community approval and active involvement in projects conducted by researchers may help to restore confidence of ethnopractitioners and provide insights into factors that underlie conservation and management of biodiversity and local ecosystems in rural settings.

3.4. Associated tick and tick-related problems

The Bukusu community appears to be well aware of livestock ticks and associated tick-borne problems. The community participants involved in the survey described to the locals the nature of ticks and the objectives of the study in the layman's language that was well understood. From their responses, it was clear that the interviewees consistently recognised the parasitic (on-host) tick stages, which were more commonly seen than the non-parasitic stages (off-host) of ticks (Tables 1a and 1b), as well as the intensity of infestations (Table 1b).

Results from Table 1b correspond to tick-related problems identified in Table 2. This confirms that every domestic animal hosts a variety of tick species. Ticks were traditionally recognised by their colour, size, type of life cycle and on-host feeding sites and habitat preference (Table 1a). A considerable proportion of local people were familiar with the locations of different ticks on livestock, including ears, udder, scrotum, tail/anal region, dewlap/neck, between legs and head. They also recognised tick and tick-related problems (T&TRPs) as serious constraints to livestock production in the study area (Table 2). The description of ticks on the host animals

Table 2Tick-borne diseases and/or related problems identified by livestock farmers as percentage of the respondents ($n = 272$).

Tick-borne diseases and/or related problems	Equivalent Bukusu name(s) used to describe tick-borne diseases and/or related problems	% respondents identifying the problems
Livestock diseases per se without specifications	<i>Bulwale bwechikhafu</i>	16.50
Anaemia	<i>Kamafuki khupungukha</i>	13.97
Poor performance of the animals	<i>Fayida yechikhafu embii</i>	12.87
Inflicted wounds and damage to skin	<i>Bulwale bwa kamakonjo khwisielo</i>	12.13
Irritation to animals	<i>Khukhwiyakala</i>	9.56
East Coast fever (ECF)	"Kamabumba"	7.72
Deaths	<i>Lifwa</i>	6.62
Weakening animals	(<i>Búúlu</i>) (<i>Butekehele</i>) <i>Khukhaywa kamani</i>	5.88
Anaplasmosis	"Kamabumba"	2.94
Heartwater	"Kamabumba"	2.57
General economic losses	<i>Asarah khulondekhana nende bulwale bwechikhafu</i>	2.57
Fever	"Walusee"	1.84
Gall sickness (disease)	–	1.25
Non-respondents	<i>Sibamanyilekhwo bulwale bwechikhafu tawe</i>	3.58

No appropriate/equivalent Bukusus name/statement describing the tick-borne disease and/or related problems.

(Table 1a) together with the problems caused by ticks (Table 2) reflected the different tick species that occurred in the study area.

Ticks are known to cause a number of serious livestock problems in the study area (Table 2). Descriptions of the ticks, including their predilection feeding sites on their host animals (Table 1a), were sufficient to confirm their taxonomic status to the genus level using voucher specimens from the Kenya National Museums (KNM) as *Rhipicephalus* spp., *Amblyomma* spp. and *Boophilus* spp. A majority of respondents classified ticks by their colours (77.1% of the responses), followed by type of life cycle (13.1% of the responses) and last by tick size (9.8% of the responses). However, classification by type of life cycle appears to be influenced by modern scientific knowledge. The highest number of responses identified the ear (25.4% of responses) as the most preferred site of tick attachment on the host animals (Table 1a), in all probability associated with the vector of East Coast fever, *Rhipicephalus appendiculatus*. A good proportion of respondents knew that ticks cause diseases to livestock (16.5%), poor productivity in affected animals (12.9%), death to livestock (6.6%), and some individuals (2.6%) were also aware of economic losses resulting from tick infestation in livestock populations (Table 2). The generation of ethnoknowledge through this type of interaction between community ethnopractitioners and conventionally trained scientists yielded valuable epidemiological information on historic and present patterns in livestock tick infestation and distribution (Hempel, 2002).

3.5. Ethnocontrol of livestock ticks in the community

While community-based traditional knowledge of tick control and management was not widely shared, it was apparent that ticks have been traditionally controlled and managed by a variety of methods (Table 3). These methods included: handpicking, on-host use of ethnobotanical suspensions (prepared from one or more of over 150 documented plants) to kill the ticks and prevent re-infestation, fumigation of infested cattle with smoke derived from burning ethnobotanical products, burning pastures, rotational grazing ethnopractices, livestock quarantine, predation by birds, cleaning cattle shed and burning residues believed to contain ticks and dusting of livestock with ash made from ethnobotanical products. These ethnotactics appear to be used in a complementary and integrated way, depending upon the level of tick infestations that resembles the modern integrated pest management (IPM) strategy. The ethnobotanicals were deployed in a number of methods that targeted a wide range of livestock tick species, either on-host or

Table 3

Frequency of the ethnomethods used by respondents in tick control and management ($n = 272$).

Ethnomethods	% respondents using the ethnomethod
Handpicking	34
Ethnobotanical products (plants and plant products)	20
Pasture spelling/grazing practices	14
Burning pastures infested with livestock ticks	14
Fumigation with smoke from burning ethnobotanical products	6
Application of kerosene, magadi soda, livestock urine, soap, fish residues, cow dung and grease (alone or mixed) on cattle	3
Livestock quarantine	3
Predation by birds	3
Cleaning cattle shed and burning residues believed to contain ticks	2
Dusting of livestock with ash from ethnobotanical products	1

Table 4

Methods of deployment of ethnobotanical products ($n = 272$).

Methods of use	% of respondents using the method
Pouring the suspension on the animal's body surface	74
Fumigation with smoke from burning ethnobotanical products	11
Rubbing livestock body with ethnobotanical products	5
Dusting the body of animals with ethnobotanical products	5
Pastures with anti-tick vegetation	2
Steaming by boiling ethnobotanical products soaked in water	1
Hanging bouquet in cattle shed to repel livestock ticks	1
Other methods	1

off-host, including pouring the suspension on the animal's body, fumigation (gaseous from ethnobotanical products directed to the animals in the cattle shed to suffocate or poison the ticks within), rubbing livestock body with ethnobotanical products, dusting livestock body with ethnobotanical products, development of tick free pastures, steaming (steam from ethnobotanical products directed to the animals in the cattle shed to suffocate or poison the ticks within) and hanging bouquet in the cattle shed to repel livestock ticks (Table 4). However, in-depth scientific studies have not been conducted to elucidate the efficacy of these ethnopractices in protecting livestock from tick disease vectors, particularly those involving the use of ethnobotanicals, which may lead to the discovery of useful anti-tick agents. These may have potential in modern integrated tick management (ITM) strategy that best fits the rural smallholder livestock farmer.

3.6. Secondary data and documented anti-tick ethnobotanicals

In total, 154 plants were identified as being used by different communities, tribes, clans and sub-clans living in Bungoma district. An extensive literature search was undertaken to document the current status of research on the plants (Appendix 1). Some of the plant species were reported in literature to be used in ethnomedicine and other cultural activities of other communities. However, no scientific literature was uncovered for a large number of plants (about 101 plant species, particularly those species belonging to the families: Lamiaceae, Flacourtiaceae, Acanthaceae and Ebenaceae) used by the communities for their apparent acaricidal and/or insecticidal effects. For some of these documented plants, literature search was extended to taxonomically related species. A large group of other plants that were repeatedly mentioned in the survey had been investigated for pesticidal, insecticidal, acaricidal or other bioactive properties and/or constituents (Appendix 1). On the other hand, some plant species (e.g. *Cleome gynandra*), that previously had been shown to have very good acaricidal properties in laboratory studies, were mentioned only once or twice by the respondents. This presented some dilemma in selecting promising candidate anti-tick plant species for detailed follow up investigations. However, based on key respondents' information, as well as acaricidal, insecticidal, pesticidal or bioactivity information from literature, eight plant species (*Tagetes minuta* L. (029-BGM-Mwi/2002), *Tithonia diversifolia* (Hemsl.) A. Gray (015-BGM-Muf/2002), *Juniperus procera* Endl. (134-BGM-Elg/2002), *Solanecio manii* (Hook. f.) C. Jeffrey. (106-BGM-Mwi/2002), *Senna didymobotrya* (Fresen.) H.S. Irwin & Barneby (132-BGM-Web/2002), *Lantana camara* L. (043-BGM-Mwi/2002), *Securidaca longepedunculata* Fres. (018-BGM-Mec/2002) and *Hoslundia opposita* Vahl.

(133-BGM-Bul/2002) were initially selected for laboratory evaluations. Of these, two plants (*Tagetes minuta* and *Tithonia diversifolia*) were chosen for in-depth detailed scientific studies in the laboratory and the field. The results of these studies will be reported elsewhere.

Additionally, some tick ethnocontrol applications that were found in the lay literature and mentioned by some respondents but that did not emerge in a scientific literature search include the use of cow dung, cattle urine, fish residues and soap. The acaricidal and behavioral effects of these on ticks and in their control would be interesting to evaluate.

4. Conclusions

The survey of indigenous tick-management knowledge of the Bukusu community revealed a wealth of preserved knowledge on plants, plant products, and anti-tick ethnopractices associated with the traditional management of ticks by rural livestock farmers. Evidence from respondents' information revealed that some of the plants used were brought from their original area/location to Bungoma by immigrants. Some plants and anti-tick ethnopractices had very few ethnoknowledge references in the literature; perhaps, they were truly indigenous to the Bukusu community or perhaps relevant references could not be accessed. Those plants thought to be indigenous to the Bukusu community and with traditionally claimed effects on livestock ticks numbered more than 101, in families such as Lamiaceae, Flacourtiaceae, Acanthaceae and Ebenaceae etc. Ritual anti-tick practices were complex and comprised burning pastures, livestock quarantine, types of grazing practices, cleaning cattle sheds, burning or burying residues of cattle sheds, bird predation, feeding animals on natural salty soils locally called "*silongo*", and applying kerosene, soap, fish residues, cowdung, cattle urine,

grease, ash, magadi soda, and sisal juices on cattle. Nevertheless, some of the local claims of acaricidal properties of the ethnobotanicals have been supported by scientific studies reported in the literature. Further elucidation of the science underlying the efficacy of these plants, plant products, and anti-tick ethnopractices may lead to the discovery of useful anti-tick agents and tactics that can be integrated in tick control and management programmes for the wellbeing of livestock industry in Africa. This may also foster wider acceptance of these ethno remedies by different stakeholders and help restore recognition accorded to ethnopractices held in the past.

Acknowledgements

The first author wishes to acknowledge International Foundation for Science (IFS) for sponsoring this study under research grant number AB/12782, and the International Centre of Insect Physiology and Ecology (ICIPE) for some financial and material support under the Dissertation Research Internship Programme (DRIP). The authors greatly appreciate the assistance of Mr. Simon G. Mathenge of the university herbarium, University of Nairobi, Kenya, and Mr. Cylus Wekesa Lantana Sinino for the identification of plant species. We are also grateful for the co-operation of Bungoma ethnopractitioners, local primary schools, Bungoma District Veterinary Officer, Bungoma District Education Officer, local administration (office of the President of the Republic of Kenya), village/clan elders and leaders, livestock farmers and Agricultural Extension Officers, throughout the study period.

Appendix 1. An enumeration of the documented plants and plant products used for the management of livestock ticks in the Bukusu community, western Kenya

Scientific name	Local and/or indigenous name(s)	Family name	Growth habit or life form	Frequency of mention for use	Plant part(s) used (application form)	Plant uses	References
<i>Acanthus pubescens</i> (Thomson ex Oliv.) Engl.	Kamarakaru	Acanthaceae	Shrub/herb	✓	Leaf/stem (ash/powder/decoction)	B, C	Baerts and Lehmann (1989, 1991)
<i>Acanthus arboreus</i> Forssk.	Kamarakaru	Acanthaceae	Shrub/herb	✓	Leaf/root (ash/powder/decoction)	C B B	Desouter (1991) Ssegawa and Kasenene (2007)
<i>Agave sisalana</i> Perrine ex Engelm.	Kamakonge or Kumukonge	Agavaceae	Forb (sub-caulescent perennials) (succulent rosette, up to 6 feet tall (1.8 m), 6 feet spread (1.8 m))	✓✓✓✓✓✓✓ ✓✓✓✓✓✓✓ ✓✓✓✓✓✓✓ ✓✓✓✓✓✓✓ ✓✓✓✓✓✓✓ ✓✓✓✓✓✓✓ ✓✓	Juice/fleshy leaves (smoking-using dried leaves; leaves and stems fermented in urine and applied; leaves dried and ground to flour and applied as dust; aqueous extract made from chopped leaves and stem and applied while still fresh)	B, I	Johns et al. (1995), Neuwinger (2000), UWMG (2003), Amer et al. (2004) Wilcox and McGeorge (1912), Schwartz et al. (1957), Faninger and Markovic-Brisk (1960), Morton (1977), Lin (1984), Nobel and Berry (1985), Ding and Chen (1989), Zullo and Azzini (1989), Pires and Purchio (1991), Chand (1991), Xu and Xu (1993), Ding and Tian (1993), FAO (1993), Adoyo et al. (1997), Chang and Chen (1997), Assefa and Williamson (2001), Schmidt (2003a)
<i>Aloe latifolia</i> Haw.	Kimitiokotioko 1	Aloeaceae	Herb/shrub	✓✓✓	Leaf/juice (aqueous)	B, T	Watt and Breyer-Brandwijk (1962), Getahun (1976), ITDG and IIRR (1996)
<i>Aloe elgonica</i> Bullock	Kimitiokotioko 2	Aloeaceae	Herb/shrub	✓	Leaf/juice (aqueous)	B, T	Watt and Breyer-Brandwijk (1962), Getahun (1976), ITDG and IIRR (1996)
<i>Ozoroa insignis</i> Del. ssp. <i>reticulata</i> (Baker f.) J.B.Gillett	Kumwandanda	Anacardiaceae	Tree	✓✓	A milky resin, bark/root/leaf	I, B, F	He et al. (2000), Araya (2001), He et al. (2002), Nyazema (2002)
<i>Rhus natalensis</i> Bernh. ex Krauss var. <i>macrocarpa</i> (Schweinf.) Cufod.	Kumusangula kumukhasi	Anacardiaceae	Tree	✓	Aerial parts/root/seed	I, F, C, B	Janet (1986), Berger (1994), Hyde et al. (2011)
<i>Rhus vulgaris</i> Meikle	Kumusangula kumusecha	Anacardiaceae	Tree	✓	Aerial parts/root/seed	I, F, C, B	Berger (1994)
<i>Lannea schimperi</i> (Hochst.ex A.Rich.) Engl.	Kumubumbu	Anacardiaceae	Tree	✓	Bark/root/leaf	F, C, B	Marin (1999), Jeruto et al. (2008)
<i>Annona senegalensis</i> Pers. ssp. <i>senegalensis</i>	Kumufwora	Annonaceae	Tree	✓	Leaf/bark/root	I, F	Stockbauer (2003)
<i>Annona chrysophylla</i> Boj.	Kumufwora	Annonaceae	Tree	✓	Leaf/bark/root	B	Toyang et al. (1995)
<i>Carisa edulis</i> Vahl.	Sirwa	Apocynaceae	Shrub	✓✓✓	Leaf/fruit/root	I, B, F F B	Stockbauer (2003) Zemedede and Mesfin (2001) Ibrahim et al. (1999), Jawaidd et al. (2011)
<i>Bidens pilosa</i> L.	Engwesi Ingwesi (Wanga)	Asteraceae	Weak annual herb	✓✓✓✓✓✓✓	Whole plant Aerial parts	I, R, Af, B, F	Chih et al. (1996), Anon. (2001d), Taylor (2002)
<i>Microglossa pyrifolia</i> (Lam.) O. Kuntze	Enguu Inguu (Wanga)	Asteraceae	Climbing shrub	✓✓✓	Whole plant-leaf-dust (used to make also suspension)	T, B	ITDG and IIRR (1996)
<i>Chrysanthemum cinerariifolium</i> (Trevir.) Vis.	Pyrethrum?	Asteraceae	Herb	✓✓✓✓✓✓✓	Flower/leaf/stem/aerial parts (used to make suspension/dust)	T, P, A, B, I	Cremlyn (1978)
<i>Tagetes minuta</i> L.	Nanjaka Nanzaka (Wanga)	Asteraceae	Herb/weed/forb	✓✓✓✓✓✓✓ ✓✓✓✓✓✓✓ ✓✓✓✓✓✓✓ ✓✓✓✓✓✓✓	Leaf/flower (dust, suspension- acqoues and hanging bouquet)	T, P, A, B, I	Getahun (1976), Graham et al. (1980), Berger (1994), Vasudevan et al. (1997), Anon. (2001d)
						F	Vasudevan et al. (1997), Leung and Foster (2003)
						I	Kennewell (1990)
						Af	Jacobson (1990)
						P	Grainge and Ahmed (1988)

Scientific name	Local and/or indigenous name(s)	Family name	Growth habit or life form	Frequency of mention for use	Plant part(s) used (application form)	Plant uses	References
<i>Solanecio mannii</i> (Hook. F.) C.Jeffrey	Nandebe (Bukusu) Mwathathia (Kikuyu) Omukhupafula (Banyala)	Asteraceae	Shrub	✓✓✓✓✓✓ ✓✓✓✓✓✓ ✓✓✓✓	Whole plant	A, C	University of Nairobi herbarium
<i>Markhamia lutea</i> (Benth.) K. Schum.	Kumusiola	Bignoniaceae	Tree	✓✓✓✓✓✓	Leaf/stem	I B C	Anon. (2003c) Kayonga and Habiyaemye (1987), New Agriculturist (2003), Rahmatullah et al. (2010), Jawaid et al. (2011) UWMG (2003)
<i>Stereospermum kunthianum</i> Cham, Sandrine Petit.	Kumutomolo (1)	Bignoniaceae	Small tree	✓✓✓✓✓✓ ✓✓✓✓✓✓ ✓✓✓✓✓✓ ✓✓✓✓✓✓ ✓✓✓✓✓✓ ✓✓✓✓✓✓	Root/bark (used to make hanging bouquet/suspension)	B, F, C, I	Abbiw (1989), Onegi et al. (2002), Petit (2003), Ching et al. (2009a,b,c,d), Compaoré et al. (2011), Jawaid et al. (2011), http://www.worldagroforestrycentre.org/sea/products/afdbases/af/asp/SpeciesInfo.asp?SpID=1556
<i>Jacaranda mimosifolia</i> D. Don	Chakaranda	Bignoniaceae	Tree	✓✓	Leaf/root/stem	C	Malerich and Trauner (2003), Rahmatullah et al. (2010) Dos Santos and Miller (1997)
<i>Piliostigma thonningii</i> (Schumach.) Milne-Redh.	Kumulamalama	Caesalpiniaceae	Perennial tree/shrub	✓✓✓✓✓✓	Leaf/root/stem (water extract) Barks/leaf/root (used to make suspension)	B T, B, C, F	Rojas et al. (2006), Rahmatullah et al. (2010), Jawaid et al. (2011) Ogundaini (1999), Oni (2001), Anon. (2001b), Le Houérou (2003)
<i>Senna siamea</i> (Lam.) Irwin & Barneby.	Mbekoraisi or Kumuraisi	Caesalpiniaceae	Tree	✓✓✓✓	Root/stem/leaf (used to make suspension)	F T, I, R	Zemede and Mesfin (2001) Adoyo et al. (1997), Mansingh and Williams (1998), Anon. (2001d)
<i>Senna singueana</i> (Del.) Lock.	Kumusilamosi	Caesalpiniaceae	Tree	✓✓✓	Fruit/root/leaf/bark (used to make suspension)	B, T, I	Adoyo et al. (1997), Mansingh and Williams (1998)
<i>Tamarindus indica</i> L.	Kumukhua	Caesalpiniaceae	Medium to large tree reaching 30 m in height.	✓✓✓✓	Root/bark/leaf (used to make suspension)	I, B	Imbabi et al. (1992), El-Siddig et al. (1999), Rao et al. (1999), Gurumurthy and Sreenivasa (2000)
<i>Senna didymobotrya</i> (Fresn.) Irwin & Barneby	Kumupinupinu	Caesalpiniaceae	Tree	✓✓✓✓✓✓ ✓✓✓✓	Root/bark/leaf (used to make suspension)	B T, B	Zemede and Mesfin (2001) Fernandes et al. (1985), Mansingh and Williams (1998), UWMG (2003)
<i>Tylosema fassoglensis</i> (Kotschy ex. Schweinf.) Torre. & Hillc.	Kumukayu	Caesalpiniaceae	Perennial herb/shrub	✓	Whole plant	T, F	Anon. (2001b)
<i>Gynandropsis gynandra</i> (L.) Briq.	Chisaka (Bukusu) Tsisaka (Wanga) Ejobyo (Luganda) Akeyo (Luo) Esaka (Banyala)	Capparidaceae	Herb	✓	Whole plant	A	Malonza et al. (1992), Dipeolu et al. (1992), Torto and Hassanali (1997), Lwande et al. (1999)
<i>Maytenus senegalensis</i> (Lam.) Exell.	Kumwayakhafu	Celastraceae	Tree	✓✓	Fruit/leaf/stem/bark/root	B T B	UWMG (2003) Ndung'u et al. (1995, 1999) El Tahir et al. (1998), El and Satti (1999), Hussein et al. (1999), Bhatt et al. (2001), Kumar (2003)

Scientific name	Local and/or indigenous name(s)	Family name	Growth habit or life form	Frequency of mention for use	Plant part(s) used (application form)	Plant uses	References
<i>Maytenus heterophylla</i> (Eckl. & Zehy) N. K. B. Robson.	Kumuwwa No. 1	Celastraceae	Tree	✓	Fruit/leaf/stem/bark/root	F, B	Working Paper FGR/18E (2001)
<i>Maytenus arbutifolius</i> (Hochst. ex A. Rich.) R. Wilczek.	Kumuwwa No. 2	Celastraceae	Tree	✓	Fruit/leaf/stem/bark/root	B, F	Getahun (1974)
<i>Terminalia mollis</i> Laws.	Kumukhonge	Combretaceae	Tree	✓✓✓	Stem/root/bark	P, T, I	Berger (1994)
<i>Combretum molle</i> R. Br. ex G. Don.	Kumukimila	Combretaceae	Tree	✓✓✓	Stem/root/bark	P, B, C, T, I	Berger (1994), Stockbauer (2003)
<i>Combretum collinum</i> Fresen.	Kumulaha	Combretaceae	Tree	✓✓✓	Resin/root/bark/leaf	P, T	Berger (1994), Odda et al. (2008)
<i>Combretum binderianum</i> Kotschy.	Kumulaha	Combretaceae	Tree	✓✓✓	Resin/root/bark/leaf	P, T	Berger (1994)
<i>Combretum elgonense</i> Exell.	Kumulaha kumukalukha	Combretaceae	Tree	✓	Resin/root/bark/leaf	P, T	Berger (1994)
<i>Vernonia amygdalina</i> Del.	Kumululusia or Kumwirurusia	Compositae	Shrub or small tree	✓✓✓✓✓✓✓	Whole plant	T, F, B, I	Getahun (1976), Toyang et al. (1995), Adoyo et al. (1997), Asfaw and Tadesse (2001)
<i>Vernonia lasiopus</i> O. Hoffm.	Namatuma	Compositae	Shrub	✓✓	Whole plant	T, F	Adoyo et al. (1997)
<i>Vernonia grantii</i> Oliv.	Endwasi	Compositae	Shrub	✓✓	Whole plant	T	Adoyo et al. (1997)
<i>Tithonia diversifolia</i> (Hemsl.) A. Gray	Kamang'ulie or Kamaua or Kiming'ulie	Compositae	Herb	✓✓✓✓	Whole plant/paste	I, P,T, B, F	Adoyo et al. (1997)
<i>Ipomoea batatas</i> (L.) Lam.	Kamabwoni (Chindabii Chekamabwoni)	Convolvulaceae	Herb	✓✓✓	Leaf/stem/sap	B	Cevallos-Casals and Cisneros-Zevallos (2002)
<i>Cupressus lusitanica</i> Mill.	Kumusayiprasi	Cupressaceae	Tree	✓✓✓✓✓✓	Leaf/stem/bark	C, I	Personal communication with herbalists
<i>Juniperus procera</i> Hochst. ex Endlicher.	Kumutarakwa (1)	Cupressaceae	Tropical African timber tree with fragrant wood	✓✓✓✓✓	Leaf/root/bark	I, B	Getahun (1976), Anon. (2001d)
<i>Euclea divinorum</i> Hiern.	Kumuchanjaasi	Ebenaceae	Tree	✓✓✓✓✓✓✓	Root/leaf/bark	I B	Hines and Eckman (1993) Homer et al. (1990), Homer et al. (1992), Dagne et al. (1993), Hines and Eckman (1993), van Grinsven et al. (1999), Lukwa et al. (2001)
<i>Euphorbia heterochroma</i> Pax.	Kumutua (1)	Euphorbiaceae	Succulent shrub	✓✓✓	Sap/stem (used to make suspension)	C T B A	Hines and Eckman (1993) Kaaya et al. (1995) Watt and Breyer-Brandwijk (1962), Philippe and Sura Faucon (2003), Schmidt (2003b) ICIPE Annual Report (1998/99), Regassaa (2000)
<i>Euphorbia tirucalli</i> L.	Kumusongofwa	Euphorbiaceae	Perennial thornless cactus-like tree	✓✓✓✓✓	Whole plant (used to make suspension)	I T B C	Hines and Eckman (1993), Schmidt (2003b), Clevinger (2003) Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Watt and Breyer-Brandwijk (1962), Getahun (1976), Duke (1983a), Hines and Eckman (1993), Bowen and Hollinger (2002) Hines and Eckman (1993)

Scientific name	Local and/or indigenous name(s)	Family name	Growth habit or life form	Frequency of mention for use	Plant part(s) used (application form)	Plant uses	References
<i>Euphorbia candelabrum</i> Kotschy.	Kumutua (2)	Euphorbiaceae	Tree	✓✓✓	Leaf/stem/sap (used to make suspension)	T A C B	Kaaya et al. (1995), Regassaa (2000) ICIPE Annual Report (1998/99) Hines and Eckman (1993) Watt and Breyer-Brandwijk (1962), Hines and Eckman (1993), Schmidt (2003b)
<i>Margaritaria discoidea</i> (Baill.) G.L.Webster	Erionoi (Teso) Atego (Luo) Kamenyabazi (Luganda) Muremampango (Runyankore)	Euphorbiaceae	Tree	✓	Whole plant (used to make suspension while wood smoke repels mosquitoes and snakes)	I T A B C	ICIPE Annual Report (1998/99), Schmidt (2003b), Adedapo et al. (2009) Regassaa (2000) Kaaya et al. (1995) Irvine (1961), Watt and Breyer-Brandwijk (1962), UWMG (2003), Schmidt (2003b) Hines and Eckman (1993), Schmidt (2003b)
<i>Bridelia scleroneura</i> Mull. Ag.	Kumunyekerwe	Euphorbiaceae	Tree/shrub	✓	Stem/root and or bark	I B F C T	Hines and Eckman (1993) Anon. (2001c), Schmidt (2003b), MTDP (2003) Persson (1986) Persson (1986) Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000)
<i>Bridelia micrantha</i> Hochst. Baill.	Kumulondamwombe or Kumukhulang'wa	Euphorbiaceae	Small tree/herb	✓✓✓✓✓✓ ✓✓✓✓	Leaf/root/bark (used to make suspension)	I T F B C S B	Hines and Eckman (1993), Anon. (2001d) Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Tredgold (1990), Hines and Eckman (1993), Asfaw and Tadesse (2001) Rulangaranga (1989), Hines and Eckman (1993), Abo and Ashidi (1999), Izelle (2002), Green et al. (2011) Hines and Eckman (1993) Tredgold (1990) Kayonga and Habiyaemye (1987), Maundu et al. (2001), Schmidt (2003b)
<i>Erythrococca bongensis</i> Pax.	Lupiriapiria	Euphorbiaceae	Tree	✓✓	Stem/root/leaf	T F B	Regassaa (2000), Kaaya et al. (1995), ICIPE Annual Report (1998/99) Anon. (1968) Schmidt (2003b)
<i>Neoboutonia melleri</i> (Müll. Arg.) Prain	Kumulongo	Euphorbiaceae	Tree/shrub	✓✓✓✓	Stem/root (used to make suspension)	T B	Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Dewees (1995), Maundu et al. (2001)
<i>Macaranga kilimandscharica</i> Pax.	Kaptebema (Sebei) Kumulongo	Euphorbiaceae	Tree Tree/shrub	✓✓✓✓✓	Stem/root (used to make suspension)	B T C	Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000), Schmidt (2003b) Dewees (1995)

Scientific name	Local and/or indigenous name(s)	Family name	Growth habit or life form	Frequency of mention for use	Plant part(s) used (application form)	Plant uses	References
<i>Croton sylvaticus</i> Hoschst. ex Krauss	Kumuchwichwi	Euphorbiaceae	Tree	✓	Seed	T B	Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Watt and Breyer-Brandwijk (1962), Mwangi et al. (1998), Schmidt (2003b) Fernandes et al. (1985), Stockbauer (2003)
<i>Croton macrostachyus</i> Hochst. ex Delile	Kumuchwichwi or Kumutotoa or Kumukunusia or Kumutoboso (Saboti)	Euphorbiaceae	Tree	✓✓	Leaf/bark/root (suspension)	I F B T C	
<i>Croton megalocarpus</i> Hutch.	Kumukunusia kumukeni or Kumutotoa kumukeni	Euphorbiaceae	Tree	✓✓	Leaf/bark/root (used to make suspension)	T C B F C T	Fernandes et al. (1985) Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Fernandes et al. (1985), Hines and Eckman (1993) Harjula (1980), Politz and Lekeleley (1988), Minja (1989), Gachathi (1989), ICRAF (1992), Hines and Eckman (1993), Kokwaro (1993), Minja (1994), Masinde (1996), ITDG and IIRR (1996), Schmidt (2003b) Hines and Eckman (1993) Hines and Eckman (1993) Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) van Rijn and Tanigoshi (1999), Mitchell and Ahmad (2006) Getahun (1976), Duke and Wain (1981), Core (1981), Fernandes et al. (1985), Toyang et al. (1995), UWMG (2003), Schmidt (2003b) Mahgoub and Ahmed (1996) Fernandes et al. (1985)
<i>Ricinus communis</i> L.	Kumuroborobo or Kumubono	Euphorbiaceae	Tree/shrub	✓✓✓✓	Seed/stem/root (used to make suspension)	A B I P	
<i>Hymenocardia acida</i> Tul.	Nangoso	Euphorbiaceae	Mostly found in savanna as woody plant/small shrubs/tree	✓	Bark/leaf (used to make decoction)	T B	Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Irvine (1961), Marin (1999), Schmidt (2003b)
<i>Drypetes gerrardii</i> Hutch.	Kumwilima	Euphorbiaceae	Tree	✓✓	Leaf/root (used to make decoction)	T B	
<i>Flueggea virosa</i> (Roxb. ex Willd.) Royle	Lubwili	Euphorbiaceae	Shrub	✓✓	Whole plant (used to make decoction)	T I	Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Schmidt (2003b) Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Anon. (2003a), Lister (2003)

Scientific name	Local and/or indigenous name(s)	Family name	Growth habit or life form	Frequency of mention for use	Plant part(s) used (application form)	Plant uses	References
<i>Sapium ellipticum</i> Hochst. ex Krauss Pax	Kumuchaso	Euphorbiaceae	Tree	✓✓✓	Root/bark (used to make decoction)	T F (Fuits) B B T	Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Asfaw and Tadesse (2001) Samuelsson et al. (1992), Schmidt (2003b) Schmidt (2003b) Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000)
<i>Phyllanthus ovalifolius</i> Forssk.	Kumusekese	Euphorbiaceae	Shrub	✓✓	Aerial parts	T B	Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Schmidt (2003b)
<i>Phyllanthus muellerianus</i> (Kuntze) Exell.	Kumutikaa	Euphorbiaceae	Tree	✓✓	Leaf	T B	Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Schmidt (2003b)
<i>Clutia richardiana</i> Moll.Arg.	Nabilikhe or Luyebeye or Lulako	Euphorbiaceae	Shrub	✓	Aerial parts (used to make concoction)	T B	Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Muhammad et al. (1999), Schmidt (2003b), Abourashed et al. (2003)
<i>Clutia abyssinica</i> Jaub. & Spach	Nabilikhe or Luyebeye or Lulako	Euphorbiaceae	Shrub	✓	Aerial parts	T B	Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Duke (2002), Schmidt (2003b)
<i>Clutia mollis</i> Pax.	Nabilikhe or Luyebeye or Lulako	Euphorbiaceae	Shrub	✓	Barks/leaves (used to make concoction)	T B	Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Schmidt (2003b)
<i>Manihot esculenta</i> Crantz.	Kumwoko	Euphorbiaceae	Perennial woody shrub	✓✓✓	Root/bark/stem (used to make concoction)	B T	Schmidt (2003b) Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Schmidt (2003b)
<i>Acalypha racemosa</i> Wall. ex Baill.	–	Euphorbiaceae	Shrub	✓✓	Root/bark/stem (used to make concoction)	B T	Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Schmidt (2003b)
<i>Dolichos kilimandscharicus</i> Harms ex Taub.	Nandwasi	Fabaceae	Herb/shrub	✓✓✓✓	Root/stem/leaf (used to make concoction)	B B, P	Kaaya et al. (1995), ICIPE Annual Report (1998/99), Regassaa (2000) Schmidt (2003b) University of Nairobi Heerbarium Library/records on voucher specimen.
<i>Garcinia buchananii</i> Bak.	Kumukhomeli	Guttiferae	Tree	✓✓✓✓✓✓✓✓	Powdered root Twig/bark/root/fruit (used to make suspension)	P Ps, B	Golob et al. (1999) Smith (1969), Duke-Elder and MacFaul (1972), Schmidt (2003c)
<i>Psorospermum febrifugum</i> Spach.	Nangoso (2)	Guttiferae	Tree	✓	Fruit/bark/stem/leaf	B	Aubréville (1936), Dalziel (1937), Irvine (1961), Cassidy et al. (1990), Schmidt (2003c)
<i>Harungana madagascariensis</i> Lam. ex Poir.	Namalasile	Guttiferae	Tree	✓	Leaf/bark/stem	B	Woodland (1997), Irvine (1961), Schmidt (2003c), The World Bank Group (2003), Erah et al. (2003)
<i>Dovyalis macrocalyx</i> (Oliv.) Warb.	Kumusongolamunwa/ Busongolomunwa (fruit)	Flacourtiaceae	A spiny shrub	✓	Whole plant (adecoction made from the crushed leaves)	F B	Maundu et al. (1999) Chifundera (1998)
<i>Tinnea aethiopica</i> Hook. f.	Lubilikhe	Lamiaceae	Shrub	✓✓✓	Leaf	T	Jembere et al. (1995)
<i>Plectranthus barbatus</i> Andr.	Ling'ulie or Kamang'ulie	Lamiaceae	Shrub	✓	Leaf	T	Jembere et al. (1995)
<i>Plectranthus lactiflorus</i> (Vatke) ined.	–	Lamiaceae	Shrub	✓	Aerial parts	T	Jembere et al. (1995)

Scientific name	Local and/or indigenous name(s)	Family name	Growth habit or life form	Frequency of mention for use	Plant part(s) used (application form)	Plant uses	References
<i>Leonotis nepetifolia</i> (L.) R. Br.	Susuni	Lamiaceae	Shrub	✓	Aerial parts	T	Jembere et al. (1995)
<i>Hoslundia opposita</i> Vahl.	Bifwofwo	Lamiaceae	Herb	✓✓✓	Root/barks	T	Jembere et al. (1995)
<i>Ocimum kilimandscharicum</i> Baker ex Gürke	Bifwofwo	Lamiaceae	Herb	✓✓	Leaf	TC	Jembere et al. (1995)
<i>Leucas calostachys</i> Oliv.	Nasusi	Lamiaceae	Herb	✓	–	T	Jembere et al. (1995)
<i>Strychnos innocua</i> Del.	Kumulende	Loganiaceae	Tree	✓✓✓✓✓✓✓	Powder of dried leaf/seed (storage pests-weevils)	P, F (fodder)	Assefa and Williamson (2001)
<i>Strychnos spinosa</i> Lam.	Kumukhubwe	Loganiaceae	Tree	✓	Leaf/bark/fruit	B Ps C, F	Hines and Eckman (1993) Stedman et al. (1942), Volfová and Patočka (2003) Wehmeyer (1966), Fox and Norwood-Young (1982), Taylor (1986), Saka and Msonthi (1994), Phillipson (1995), Mizrahi et al. (2002), Kristensen and Lykke (2003) Puckhaber et al. (2002)
<i>Hibiscus calyphyllus</i> Cav.	Kumwarakumba	Malvaceae	Shrub	✓	Leaf/stem/aerial parts/root	F, N	
<i>Ekebergia capensis</i> Sparm.	Kumusilisi (1)	Meliaceae	Evergreen or semi-deciduous, medium-sized to large tree.	✓✓✓	Root/bark/fruit/flower/leaf/stem	T, B, C, F	World Agroforestry Centre (2003a)
<i>Ekebergia rueppeliana</i> (Fresen.) A. Rich.	Kumusilisi (2)	Meliaceae	Evergreen or semi-deciduous tree	✓✓✓	Root/bark/fruit/flowers	T, C, F, B	World Agroforestry Centre (2003b)
<i>Turraea holstii</i> Gürke	Nawili	Meliaceae	Tree	✓✓	Whole plant	B T	Mulholland et al. (1998), Rajab et al. (1998) Lindsay and Kaufman (1988), Cabral et al. (1996), Mansingh and Williams (1998), Ndumu et al. (1999), Abdel-Shafy and Zayed (2002), Borges et al. (2003)
<i>Turraea robusta</i> Gürke	Nabili	Meliaceae	Small tree	✓✓✓✓	Whole plant	T, B	Rajab et al. (1988), Bentley et al. (1992), Rajab et al. (1998), Gathirwa et al. (2008)
<i>Azadirachta indica</i> A. Juss.	Kumwarubaine (2)	Meliaceae	Tree	✓✓✓✓✓✓ ✓✓✓✓✓✓ ✓✓✓✓✓✓✓✓	Stem/bark/fruit/leaf/root (suspension/dusting/rubbing bolus/paste)	I A	ITDG and IIRR (1996), ICIPE Annual Report (1998/99), Mansingh and Williams (1998), Mulla and Su (1999), Ndumu et al. (1999), Anon. (2001d), Abdel-Shafy and Zayed (2002), Singh (2002) Lindsay and Kaufman (1988), Williams (1993), ITDG and IIRR (1996), ICIPE Annual Report (1998/99), Ndumu et al. (1999), Wilson and Mansingh (2002), Blair and Mansingh (2002), Abdel-Shafy and Zayed (2002), Mansingh and Williams (2002)

Scientific name	Local and/or indigenous name(s)	Family name	Growth habit or life form	Frequency of mention for use	Plant part(s) used (application form)	Plant uses	References
<i>Melia azedarach</i> L.	Kumwarubaine (1)	Meliaceae	Tree	✓✓✓✓✓✓ ✓✓✓✓✓✓ ✓✓✓✓✓✓✓✓	Stem/bark/fruit leaf/root (used to make suspension/dusting/rubbing)	T	Lindsay and Kaufman (1988), Cabral et al. (1996), Mansingh and Williams (1998), Borges et al. (2003)
						P	Isman et al. (1996), Brechelt (2002), Singh (2002)
						B	UWMG (2003), Thapa and Pathak (2002)
						Af	Vellaikumar (1997), ICIPE Annual Report (1998/99)
						T	Williams (1993), Ndumu et al. (1999), Anon. (2001d), Abdel-Shafy and Zayed (2002)
						A	Lindsay and Kaufman (1988), Cabral et al. (1996), Mansingh and Williams (1998), Borges et al. (2003)
						P	Stoll (1988), Neupane (1992), Berger (1994), Kroschel (1996), Batchner (2000)
						Af	Chen et al. (1996), Nardo et al. (1997), Vellaikumar (1997)
						B	Andrei et al. (1990), Zakir et al. (1991), Villamil et al. (1995), HerbWeb (2000)
<i>Acacia macrothyrsa</i> Harms.	Kumukholondo	Mimosaceae	Shrub/tree	✓✓✓	Root/leaf/bark (used to make hanging bouquet/suspension)	B, C, T, F	Al-Sharook et al. (1991), Arias et al. (1992), Isman et al. (1996), Cabral et al. (1996), Nardo et al. (1997), Valladares et al. (1997), Adoyo et al. (1997), Schmidt et al. (1998), Singh (2002) Marin (1999), Le Houérou (2003)
<i>Albizia coriaria</i> Welw. ex Oliv.	Kumupeli (Bukusu)	Mimosaceae	Perennial tree	✓✓✓✓✓✓	Stem/root/bark/leaf	T, B	Kokwaro (1993), ITDG and IIRR (1996), Schmidt (2003d)
<i>Albizia grandibracteata</i> Taub.	Kumufunjamwe/Kumulongo/Kumunianyuni	Mimosaceae	Perennial tree	✓	Bark/leaf	T, N	–
<i>Entada abyssinica</i> A.Rich.	Kumusembe	Mimosaceae	Perennial tree	✓✓	Root/bud/leaf/ Bark (used to make suspension)	T, C, F, B	Hines and Eckman (1993), Toyang et al. (1995), Freiburghaus et al. (1998), Duke (2002)
<i>Acacia sieberiana</i> DC.	Kumulemba	Mimosaceae	Perennial Tree	✓✓✓✓	Root	F, C, B, T	Anon. (2001a)
<i>Acacia nilotica</i> (L.) Bel.	Kumulemba	Mimosaceae	Tree	✓✓✓✓	Bark/root	B, I, F B, P	Duke (1983b), Anon. (2001d) Heine and Heine (1988), Heine and König (1988), Timberlake (1987)
<i>Ficus sycomorus</i> L.	Kumukhuyu	Moraceae	Tree	✓✓✓✓✓✓	Leaf/fruit/bark/latex	T, B, F, C	Puyvelde (van) et al. (1985), Hines and Eckman (1993), Regassaa (2000)
<i>Ficus vasta</i> Forssk.	Kumutoto kumutotoa	Moraceae	Tree	✓✓✓✓	Root/fruit/bud/latex (used to make suspension)	T, F, C	Puyvelde (van) et al. (1985), Regassaa (2000)
<i>Ficus amadiensis</i> De Wild.	Kumutoto kumukobianiyuni	Moraceae	Tree	✓✓	Root/fruit/bark/latex	T, C	Puyvelde (van) et al. (1985), Regassaa (2000)
<i>Ficus thonningii</i> Blume.	Kumutoto kumusecha	Moraceae	Tree	✓	Root/fruit/bark/latex	T, F, B, C	Puyvelde (van) et al. (1985), Hines and Eckman (1993), Regassaa (2000), Stockbauer (2003)
<i>Ficus glumosa</i> Del.	Kumusilisisi	Moraceae	Tree	✓	Root/fruit/bark/latex	–	–
<i>Ficus sur</i> Forssk.	Kumukhuyu nandere	Moraceae	Tree	✓✓	Root/fruit/bark/latex (used to make suspension)	T, F, C	Puyvelde (van) et al. (1985), Regassaa (2000)

Scientific name	Local and/or indigenous name(s)	Family name	Growth habit or life form	Frequency of mention for use	Plant part(s) used (application form)	Plant uses	References
<i>Syzygium guineense</i> (Willd.) DC. ssp. <i>guineense</i>	Kumusitole kumusecha	Mrytaceae	Tree	✓	Bark/root/fruit	C, F, B	Hines and Eckman (1993)
<i>Syzygium cordatum</i> Hochst.ex C.Krauss.	Kumusemwa	Mrytaceae	Medium-large tree.	✓	Stem/root/bark/fruit	F	Trade Winds Fruit (2003)
<i>Syzygium cumini</i> L.	Kumusemwa	Mrytaceae	Medium-large tree.	✓	Stem/root/bark/fruit	F	Trade Winds Fruit (2003)
<i>Maesa lanceolata</i> Forssk.	Namaru	Myrsinaceae	Tree	✓✓	–	T, B	Getahun (1976)
<i>Lophira alata</i> Banks ex Gaertn.	Oteng (Luo)	Ochnaceae	Tree	✓	Whole plant	B	Anon. (1977)
<i>Ximenia americana</i> L.	Kumutuli	Olacaceae	Woody shrub or small tree	✓✓✓	Leaf/bark/root/fruit	P, B, C, F	Hines and Eckman (1993)
<i>Olea capensis</i> L.	Kumutamaywa	Oleaceae	Tree	✓	Leaf/bark/fruit	B F, C	Anon. (2003b) Hines and Eckman (1993), Mutangah (1996)
<i>Olea europaea</i> (L.) ssp. <i>africana</i> (Mill.) P. Green.	Kumunyubuti	Oleaceae	Woody perennial tree	✓	Bark/stem/leaf	B C	Hines and Eckman (1993), Stockbauer (2003) Palgrave (1988), Hines and Eckman (1993)
<i>Sesbania sesban</i> (L.) Merrill.	Chisubasubi (1)	Papilionaceae	Tree	✓✓✓	Leaf/stem (used to make suspension)	F B	Hines and Eckman (1993) Botanical South Africa (2003), Viable Herbal Solutions (2003)
<i>Kotschyia africana</i> Endl.var. <i>bequaertii</i> (De Wild) Verdc.	Chisubasubi (2)	Papilionaceae	Tree	✓	Various parts of the plant	T	Matzigkeit (1990)
<i>Sesbania macrantha</i> Welw. ex Phillips and Hutch.	Chisubasubi (3)	Papilionaceae	Tree	✓✓✓	Leaf/stem (used to make suspension)	T	Matzigkeit (1990)
<i>Indigofera emarginella</i> Steud. ex A. Rich.	Lusisakwe	Papilionaceae	Shrub	✓✓✓	Root/stem	T	Matzigkeit (1990)
<i>Erythrina abyssinica</i> Lam. ex DC.	Kumurukuru or Kumurembe	Papilionaceae	Perennial shrub/tree	✓✓	Leaf/bark (used to make hanging bouquet/dusting/fumigation)	T B C	Matzigkeit (1990), Mansingh and Williams (1998) Rulangaranga (1989), Hines and Eckman (1993), ITDG and IIRR (1996) Rulangaranga (1989), Hines and Eckman (1993)
<i>Tephrosia vogelii</i> Hook F.	Kumufila or Esumu	Papilionaceae	Shrub	✓✓✓✓✓✓ ✓✓✓✓✓✓ ✓✓✓✓✓✓✓✓	Leaf/stem/fruit/bud/bark/seed/root (whole plant)	A, T I Af B P T, B	Lemmens and Fryer (1917), Cremllyn (1978), Matzigkeit (1990), Toyang et al. (1995), Kambewa et al. (1997) Kambewa et al. (1997) Niang (1987), Bertram (1988) Niang (1987), Getahun (1976) Matzigkeit (1990), Cremllyn (1978) Matzigkeit (1990), William (1999)
<i>Rhynchosia resinosa</i> (A. Rich.) Baker	Kumusiuli	Papilionaceae	Tree	✓	Leaf/stem/root/bark	T, B	Matzigkeit (1990), William (1999)
<i>Macuna poggei</i> Taub.var. <i>pesa</i> (De Wild) Verdc.	Kumuchuli or Kumusiuli	Papilionaceae	Perennial climbing herb or shrub	✓	Bark/root	T	Matzigkeit (1990)
<i>Mucuna poggei</i> Taubert var. <i>poggei</i> (De Wild) Verdc.	Kumuchuli or Kumusiuli	Papilionaceae	Perennial climbing herb or shrub	✓	Aerial parts	T	Matzigkeit (1990)

Scientific name	Local and/or indigenous name(s)	Family name	Growth habit or life form	Frequency of mention for use	Plant part(s) used (application form)	Plant uses	References
<i>Mondia whytei</i> (Hook f.) Skeels	Kumukombela	Periplocaceae	Herb	✓	Root/stem?	B	Okwemba (2002)
<i>Pittosporum viridiflorum</i> Sims.	Nambaa (1)	Pittosporaceae	Shrub	✓✓	Stem bark/root/leaf	F, B	Coetzee et al. (1999), Berger et al. (2002), Seo (2002)
<i>Sorghum bicolor</i> (L.) Moench.	Kamayemba (Bukusus) Amabele (Wanga) Ammabelee (Banyala)	Poaceae	Grass	✓	Whole plant (particularly grains, leaf and stem)	I, F, B	Grieve (1931), Watt and Breyer-Brandwijk (1962), Duke and Wain (1981), Morton (1981), Perry (1980), Duke (1983c), Anon. (2001d)
<i>Saccharum officinarum</i> L.	Kimiba (Bukusu) Emikhonye (Wanga)	Poaceae	Grass	✓	Leaf/stem/juice	F	–
<i>Zea mays</i> L.	Kamayindi (Bukusu) Amatumwa (Wanga)	Poaceae	Grass	✓✓✓	Leaf/stem	B	Lukwa et al. (2001), Ramos-Escudero et al. (2011)
<i>Podocarpus falcatus</i> (Thunb.) R.Br. ex Mirb.	Kumutarakwa (2)	Podocarpaceae	Shrub	✓	Leaf/root/bark	I	Stockbauer (2003)
<i>Podocarpus latifolius</i> (Thunb.) R.Br. ex Mirb.	Kumutarakwa (3)	Podocarpaceae	Tree	✓✓	Stem/root/leaf	C	Stockbauer (2003)
<i>Securidaca longipedunculata</i> Fres.	Kumunyakasia	Polygalaceae	Tree	✓✓	Whole plant	I, C, B, F, Ps	Rukangira (2001), Jayasekara et al. (2002)
<i>Zizyphus abyssinica</i> Hochst. ex A. Rich.	Kumukomboti (1)/Kumwikalangwe	Rhamnaceae	Tree	✓✓✓	Root/bark	F	Hentgen (1985)
<i>Zizyphus mucronata</i> Willd.	Kumukomboti (2)/Kumwikalangwe/Tirokwo (Pokot)	Rhamnaceae	Tree	✓✓✓	Root/bark	F	Hentgen (1985)
<i>Prunus africana</i> (Hook.f.) Kalkman.	Kumuturu/Kumwilima	Rosaceae	Tree	✓✓✓	Bark/leaf/root	P, I, B	Cunningham and Mbenkum (1993), Schippmann (2001), World Agroforestry Centre (2003b), USDA, ARS, National Genetic Resources Program (2004)
<i>Gardenia ternifolia</i> Schumach. & Thonn.	Siuna	Rubiaceae	Shrub	✓✓✓✓✓✓ ✓✓✓✓✓✓✓✓	Root/fruit	F	Asfaw and Tadesse (2001)
<i>Pavetta oliveriana</i> Hiern.	Kumukokhakokhe	Rubiaceae	Tree/shrub	✓	Root/leaf	C, I	Fernandes et al. (1985)
<i>Pavetta crassipes</i> K. Schum.	Kumusindusianjofu/ Kumupepenambusi kumukhasi	Rubiaceae	Tree	✓✓	Root/leaf	– F, B	N Sanon et al. (2003).
<i>Teclea nobilis</i> Del.	Kumutare	Rutaceae	Tree	✓	Root/stem/bark/leaf	C, B	Hines and Eckman (1993), Marin (1999)
<i>Paullinia pinnata</i> L.	Kumukoyakoye kumulaasi	Sapindaceae	Lianas	✓	The root and root bark are applied for rubefacient purposes.	B, F	Killip and Smith (1935), Fanshawe (1948, 1953), Watt and Breyer-Brandwijk (1962), Abourashad et al. (1999), van Andel (2000), Onge (2002), Duke and Bogenschutz (2003)
<i>Harrisonia abyssinica</i> Oliv.	Sibondwe	Simaroubaceae	Tree	✓✓	Leaf/root/bark/stem	B	Bally (1937), Watt and Breyer-Brandwijk (1962), Johns et al. (1990), Kokwaro (1993), Fabry et al. (1996)
<i>Withania somnifera</i> (L.) Dunal	Nambaa (2)	Solanaceae	A perennial herb	✓✓	Stem/root/leaf	B	Pande (1999), Singh and Kumar (1998), Scartezini and Speroni (2000), McIntyre (2002), Chopra et al. (2004), Van Der Hoof et al. (2005), Cooley et al. (2009), Mirjalili et al. (2009), Ven Murthy et al. (2010), Ahmad et al. (2010)

Scientific name	Local and/or indigenous name(s)	Family name	Growth habit or life form	Frequency of mention for use	Plant part(s) used (application form)	Plant uses	References
<i>Nicotiana tabacum</i> L.	Eraba	Solanaceae	Herb	✓✓✓✓✓	Leaf/stem-dust (used to make suspension smoke and hanging bouquet)	P, A, I, B, C	Cremlyn (1978), Stoll (1988), Matzigkeit (1990), Juliette de Ba Levy (1991), Dipeolu and Ndungu (1991), Berger (1994), Mwangi (1996), ITDG and IIRR (1996), Adoyo et al. (1997), Mansingh and Williams (1998), Adewusi and Afolayan (2010)
<i>Capsicum annum</i> L.	Pilipili 1	Solanaceae	Shrub	✓✓✓✓✓ ✓✓✓✓✓ ✓✓✓✓✓	Fruit/leaf/stem (suspension, dust, smoking and hanging bouquet)	P, I, T, B, F	Stoll (1988), Berger (1994), Adoyo et al. (1997), Mansingh and Williams (1998), Regassaa (2000)
<i>Capsicum frutescens</i> L.	Pilipili 2	Solanaceae	Shrub	✓✓✓✓✓ ✓✓✓✓✓ ✓✓✓✓✓ ✓✓✓✓✓	Fruit/leaf/stem (suspension, dust, smoking and hanging bouquet)	P, I, T, B, F	Stoll (1988), Berger (1994), Adoyo et al. (1997), Mansingh and Williams (1998), Regassaa (2000)
<i>Solanum incanum</i> L.	Chindurandura	Solanaceae	Shrub	✓✓✓✓✓ ✓✓✓✓✓	Root, fruit-juice (suspension)	B, C, F	Hines and Eckman (1993), Lukwa et al. (2001), Jawaide et al. (2011), Adewusi and Afolayan (2010)
<i>Physalis peruviana</i> L.	Embunwe Emilwa (Wanga)	Solanaceae	Herb	✓✓	Stem/root/fruit/leaf (heated leaves are applied as poultices on inflammations and leaf infusion as an enema to relieve abdominal ailments in children).	F, B, C	Morton (1987), Barbadine (2003), Wu et al. (2005), Wu et al. (2006), Franco et al. (2007), Arun and Asha (2007), Pardo et al. (2008), Pinto et al. (2009), Lan et al. (2009), Wu et al. (2009), Kolar and Malbeck (2009), Yen et al. (2010)
<i>Grewia trichocarpa</i> Hochst. ex A. Rich.	Kumusitati or Lusitati	Tiliaceae	Tree	✓	Root	F, B	Brink (2009)
<i>Grewia bicolor</i> A. Juss.	Kumusitati or Lusitati	Tiliaceae	Tree	✓	Root/bark	B	http://www.pbs.org/wnet/africa/explore/sahel/sahel.vegetation.io.html (retrieved 19.08.004), Orwa et al. (2009)
<i>Steganotaenia araliaceae</i> Hochst.	Kumutomolo (1) or Kumupepenambusi kumusecha	Umbelliferae	Tree	✓	Leaf/stem/bark	B, P	Rukangira (2003)
<i>Clerodendrum myricoides</i> (Hochst.) R. Br. & Vatke.	Nangoso (3)	Verbenaceae	Tree	✓✓✓	Leaf, juice, roots	B, P, I, A B, P	Baerts and Lehmann (1991), Kasonia et al. (1991) Staner and Boutique (1937), Watt and Breyer-Brandwijk (1962), Glover et al. (1966), Lemordant (1971), Van Puyvelde et al. (1977), Lindsay and Hepper (1978), Harjula (1980), Gelfand et al. (1985), Kayonga and Habiyaemye (1987), Heine and Heine (1988), Baerts and Lehmann (1989), Gachathi (1989), Polygenis-Bigendako (1990), Kokwaro (1993), Tadesse (1994)
<i>Rotheca myricoides</i> (Hochst.) Steane & Mabb.	Nangoso (1)	Verbenaceae	Shrub	✓✓	Whole plant	B, T, B	http://members.lycos.co.uk/ethiopianplants/shinasha.html , Getahun (1976), Kebenei et al. (2004)
<i>Lantana camara</i> L.	Mukhekhe (Bukusu) Lantana (Wanga)	Verbenaceae	Herb/shrub	✓✓	Whole plant (pour on or steaming)	B, F, C, I	Adebayo et al. (1999)
<i>Lippia grandifolia</i> Hochst. ex A. Rich. var. longipedunculata Moldenke	Lukumakuma	Verbenaceae	Herb	✓✓✓✓✓	Root/leaf/stem	T ¹	Kunle et al. (2003), Mwangi et al. (1989, 1991a,b, 1994, 1995), Koumaglou et al. (1996a,b), Garneau et al. (1995), Mwangi et al. (1998)

Scientific name	Local and/or indigenous name(s)	Family name	Growth habit or life form	Frequency of mention for use	Plant part(s) used (application form)	Plant uses	References
<i>Vitex doniana</i> Sweet	Kumufutu or Kumufututu	Verbenaceae	Tree	✓✓✓✓	Bark/root/leaf/fruit (used to make smoke and hanging bouquet)	I, B, F, C	FAO (1983), RSCU (1992), Hines and Eckman (1993), Olusola et al. (1997), Rukangira (2001), Ky (2008)
<i>Vitex fischeri</i> Gürke	Kumufutumbwe	Verbenaceae	Tree	✓✓✓✓✓✓✓	Whole plant (used to make smoke and hanging bouquet)	I, F, B	Thijssen (2008)
<i>Cyphostemma adenocaula</i> (Steud. ex A. Rich.) Desc.	Kumukoyakoye	Vitaceae	Lianas	✓	Whole plant	B	Adamantidis (1956), Polygenis-Bigendako (1990), Kokwaro (1993)
<i>Cyphostemma heterotrichum</i> (Gilg & R.E. Fr.) Desc. ex Wild & R.B. Dru.	Libombwe	Vitaceae	Lianas	✓	Whole plant	B	N
<i>Rhoicissus tridentata</i> (L.f.) Wild & R.B. Drumm.	Kumukoyakoye	Vitaceae	Lianas	✓✓✓✓✓✓ ✓✓✓✓✓✓✓	Root or tuberous rootstock (lignotuber)	F, C, B	Kayonga and Habiaremye (1987), Cunningham (1990, 1995), Maundu et al. (2001), Michael (2003), Brookes and Katsoulis (2006), Gruber and O'Brien (2010)
<i>Rhoicissus revoilii</i> Planch.	Kumukoyakoye	Vitaceae	Climber	✓✓✓✓	Whole plant	B	ITDG and IIRR (1996)
<i>Cyphostemma orondo</i> (Gilg & M.Brandt) Desc.	Kumukoyakoye	Vitaceae	Lianas	✓	Whole plant	–	N
Other anti-tick ethnoagents (claimed by the Bukusu community to work best in mixed forms as agents to control and manage livestock ticks)							
	Bar soap (toxic?)						
	Fish (odours) (repellent?)						
	Cattle urine + sisal (toxic and repellent?)						
	Kerosene (toxic and repellent?)						
	Magadi soda (toxic)						Dipeolu and Ndungu (1991)
	Grease (toxic and repellent?)						
	Ash (toxic and repellent?)						
	Feeding animals on natural salty soils locally called “silongo” (repellent/toxic?)						Dipeolu and Ndungu (1991)
	Cattle cowdung (repellent)						Sika (1996), Wanzala et al. (2004)

✓, refers to a mention of a plant for its use in tick control and/or control of other livestock pests; A, plants that were found in literature to demonstrate acaricidal activities/compounds; B, plants that were found in literature to have bioactive and toxic compounds other than pesticidal/acaricidal/insecticidal constituents; P, plants that were found in literature to demonstrate pesticidal activities/compounds other than insecticidal constituents; I, plants found in literature to be resistant to attack by insects or found to demonstrate insecticidal activities/compounds; T, taxonomic affinity to plant species (at genus and family levels) known to possess bioactive, insecticidal, acaricidal activities/compounds; N, no leading bioactive, insecticidal and or acaricidal information from literature; F, human food plants, unless otherwise stated; Af, plants with antifeedant activities against insects; C, Plants with cultural implication and applications; –, information not available for documentation; S, sericulture – the cultivation of silk worms for the production of natural silk fiber; Ps, poisonous plants; ?, no consistent information obtained on the plants from field responses and different literature sources.

References

- Abbiw, D., 1989. Useful plants of Ghana. University of Ghana, Legon, Royal Botanic Gardens, Kew, London. Annotated bibliography. In: Falconer, J., Koppell, S.R.C. (Eds.), *The Major Significance of 'Minor' Forest Products: The Local Use and Value of Forests in the West African Humid Forest Zone*, Food and Agriculture Organization of the United Nations, Rome, 1990. <http://www.fao.org/docrep/t9450e/t9450e0t.htm> (12.12.11).
- Abdel-Shafy, S., Zayed, A.A., 2002. In vitro acaricidal effect of plant extract of neem seed oil (*Azadirachta indica*) on egg, immature, and adult stages of *Hyalomma anatolicum excavatum* (Ixodoidea: Ixodidae). *Veterinary Parasitology* 106, 89–96.
- Abegaz, B.S., Demissew, S., 1998. Indigenous African useful plants with special emphasis on medicinal plants and issues associated with their management. A paper presented at a regional workshop on Africa's Natural Resources Conservation and Management Surveys, 2nd to 5th March, 1998, Accra, Ghana.
- Abo, K.A., Ashidi, J.S., 1999. Antimicrobial screening of *Bridelia micrantha*, *Alchornea cordifolia* and *Boerhavia diffusa*. *African Journal of Medicine and Medical Sciences* 28, 167–169.
- Abourashad, E., Toyang, N.J., Choinski Jr., J.S., Khan, I., 1999. Two new flavone glycosides from *Paullinia pinnata*. *Journal of Natural Products* 62, 1179–1181.
- Abourashed, E.A., Ganzeria, M., Khan, I.A., Khan, S., Mossa, J.S., El-Ferally, F.S., 2003. HPLC fingerprinting and estimation of the bioactive components of *Clusia richardiana* L. as a potential hypoglycemic herbal tea. *Phytotherapy Research* 17, 657–660.
- Adamantidis, A., 1956. Monographie pastorale du Rwanda-Urundi. *Bulletin Agricole du Congo Belge* 47, 632–635.
- Adebayo, A.T., Gbolade, T.A., Olaifa, A.A., 1999. Comparative study of toxicity of some essential oils to larvae of three mosquito species. *Nigerian Journal of Natural Products and Medicine* 3, <http://www.inasp.info/ajol/journals/njnpm/vol3abs.html>. October 20, 2003.
- Adebadjo, A.A., Sofidiya, M.O., Afolayan, A.J., 2009. Anti-inflammatory and analgesic activities of the aqueous extracts of *Margaritaria discoidea* (Euphorbiaceae) stem bark in experimental animal models. *Revista De Biología Tropical* 57, 1193–1200.
- Adewusi, A.E., Afolayan, J.A., 2010. A review of natural products with hepatoprotective activity. *Journal of Medicinal Plants Research* 4, 1318–1334.
- Adoyo, F., Mukalama, J.B., Enyola, M., 1997. Kenya Woodfuel and Agroforestry Programme (KWAP), Busia, Kenya. Using tithonia concoctions for termite control in Busia District, Kenya. *ILEIA Newsletter* 13, 24, <http://www.oneworld.org/ileia/newsletters/13-4/13-4-24.htm#top> (09.10.03).
- Agyare, C., Lechtenberg, M., Deters, A., Peterleit, F., Hensel, A., 2011. Ellagitannins from *Phyllanthus muellerianus* (Kuntze) Exell.: Geraniin and furosin stimulate cellular activity, differentiation and collagen synthesis of human skin keratinocytes and dermal fibroblasts. *Phytomedicine* 18, 617–624.
- Ahmad, M.K., Mahdi, A.A., Shukla, K.K., Islam, N., Rajender, S., Madhukar, D., Shankhwar, S.N., Ahmad, S., 2010. *Withania somnifera* improves semen quality by regulating reproductive hormone levels and oxidative stress in seminal plasma of infertile males. *Fertility and Sterility* 94, 989–996.
- Alghali, M.A., 1992. Component technology development: an integrated tick management approach for livestock production in Africa. *Insect Science and Its Application* 13, 609–612.
- Al-Sharook, Z.K., Balan, Y., Jiang, Y., Hein, Z., 1991. Insect growth inhibitors from two tropical Meliaceae. Effect of crude seed extracts on mosquito larvae. *Journal of Applied Entomology* 111, 425–430.
- Amer, M.E., Abou-Shoera, M.I., Abdel-Kader, M.S., El-Shaibany, M.S., Amina, A.M.S., Abdel-Salama, A.N., 2004. Alkaloids and flavone acyl glycosides from *Acanthus arboreus*. *Journal of the Brazilian Chemical Society* 15, 1–5.
- Andrei, G.M., Coulombie, F.C., Courreges, M.C., DeTorres, R.A., Coto, C.E., 1990. Meliacine, an antiviral compound from *Melia azedarach* L., inhibits interferon production. *Journal of Interferon Research* 10, 469–476.
- Anon., 1968. Food Composition Table for Use in Africa. U.S. Department of Health, Education and Welfare, Public Health Service, Health Services and Mental Health Administration, National Center for Chronic Disease Control, Nutrition Program Bethesda, Maryland 20014 and Food Consumption and Planning Branch, Nutrition Division, Food and Agriculture Organization of the United Nations, Rome, Italy. <http://www.fao.org/docrep/003/X6877E/X6877E00.htm#TOC> (27.09.03).
- Anon., 1977. Posterior paralysis among sows kept on sawdust from the Azobe tree (*Lophira alata* var. *procera*). *Tijdschr Diergeneesk* 102, 582.
- Anon., 2001. *Acacia sieberiana* DC. <http://www.ildis.org/LegumeWeb/6.00/taxa/373.shtml> (20.10.03).
- Anon., 2001. *Tylosema fassoglensis* (Schweinf.) Torre and Hillc. <http://www.ildis.org/LegumeWeb/6.00/taxa/373.shtml> (20.10.03).
- Anon., 2001. Structure and Data on Anthriscin, Deoxydophyllotoxin, A Natural Product with Anticancer Activity Isolated from *Bridelia scleroneura* [N1949, Q67B205] Korean. Molecular Targets Development Program. Center for Cancer Research at the National Cancer Institute. <http://home.ncicfcr.gov/mtdp/compounds/403148.html> (27.09.03).
- Anon., 2001. Termite Control HDRA – The Organic Organization. Ryton Organic Gardens, COVENTRY CV8 3LG, United Kingdom. www.hdra.org.uk (27.10.01).
- Anon., 2003. *Flueggea virosa* (Roxb. ex Willd.) Voigt. Hong Kong Flora and Vegetation. <http://www.hkflora.com/index.htm> (01.10.03).
- Anon., 2003. Management Options for the Golden Apple Snail. <http://www.applesnail.net/pestalert> (12.10.03).
- Anon., 2003. On-farm Technology Testing. Sustainable Agriculture Extension Manual. UNDP SANE (Sustainable Agriculture Networking and Extension) Program, Berkeley, California. <http://www.iirr.org/saem/contents.htm#top> (10.10.03).
- Araya, E.E., 2001. State of Forest genetic Resources in Eritrea. Sub-regional Workshop FAO/IPGRI/ICRAF on the Conservation, Management, Sustainable Utilization and Enhancement of Forest Genetic Resources in Sahelian and North-Sudan Africa (Ouagadougou, Burkina Faso, 22–24 September 1998). Forest Genetic Resources Working Papers, Working Paper FGR/23E. Forestry Department, FAO, Rome, Italy.
- Arias, A.R., Schmeda-Hirschmann, G., Falcão, A., 1992. Feeding deterrence and insecticidal effects of plant extracts on *Lutzomyia longipalpis*. *Phytotherapy Research* 6, 64–67.
- Arun, M., Asha, V.V., 2007. Preliminary studies on antihepatotoxic effect of *Physalis peruviana* Linn. (Solanaceae) against carbon tetrachloride induced acute liver injury in rats. *Journal of Ethnopharmacology* 111, 110–114.
- Asfaw, Z., Tadesse, M., 2001. Prospects for sustainable use and development of wild food plants in Ethiopia. *Economic Botany* 55, 47–62.
- Assefa, F., Williamson, S., 2001. Ethiopian farmers test alternatives to pesticides. Integrated Pest Management-Farmers Field School Project. *Pesticides News* No. 52, pp. 8–9. <http://www.pan-uk.org/pestnews/pn52/pn52p8.htm> (09.10.03).
- Aubréville, A., 1936. La Flore Forestière de la Côte d'Ivoire. Paris, France.
- Baerts, M., Lehmann, J., 1989. Guérisseurs et plantes médicinales de la région des crêtes Zaire-Nil au Burundi. Musée royal de l'Afrique centrale, Tervuren, Belgique. *Annals of Science Ecology* 18, 214.
- Baerts, M., Lehmann, J., 1991. Plantes médicinales vétérinaires de la région des crêtes Zaire-Nil au Burundi. Musée royal de l'Afrique centrale, Tervuren. *Annals of Science Ecology* 21, 133.
- Baldwin, S., Cervinkas, J., 1993. Community participation in research. Proceedings of a workshop held in Nairobi, Kenya, 23–27 September 1991. Co-sponsored by the International Development Research Centre (IDRC) and the University of Nairobi, IDRC Publications, Nairobi, Kenya.
- Bally, P.R.O., 1937. Native medicinal and poisonous plants of East Africa. *Kew Bulletin* 1, 10–25.
- Barbadine, 2003. Graines et semences tropicales/Tropical seeds. http://www.barbadine.com/pages/tropical_seeds.htm (03.11.03).
- Batcher, S.M., 2000. Element stewardship abstract for *Melia azedarach* Chinaberry, Umbrella tree. In: Mandy, T., John, M., Randall, J.M. (Eds.), *The Nature Conservancy's Wild Land Invasive Species Team*, 124 Robbins Hall. Department of Vegetable Crops and Weed Science, University of California, Davis, USA, <http://tncweeds.ucdavis.edu/esadocs/documnts/meliazee.html> (10.10.03).
- Bentley, M.D., Gaul, F., Rajab, S.M., Hassanali, A., 1992. Tetranortriterpenes from *Turraea robusta* (Gürke). *Journal of Natural Products* 55, 84–87.
- Berger, A., 1994. Using natural pesticides: current and future perspectives. A report for the Plant Protection Improvement Programme in Botswana, Zambia and Tanzania. Department of Entomology, Swedish University of Agricultural Sciences, P.O. Box 7044, S-750 07 Uppsala, Sweden. Number (ISBN, ISSN): ISSN: 1104-6422. http://www.vaxteko.nu/html/sli/slu/rapport_vaxtsk.vetensk/RVX02/RVX02.HTM (12.10.03).
- Berger, J.M., Hoch, J., Neddermann, K.M., Bursuker, I., Mamber, S.W., Kingston, D.G.I., 2002. A new triterpene saponin from *Pittosporum viridiflorum* from the Madagascar rainforest. *Journal of Natural Products* 65, 11–15.
- Bertram, T.M., 1988. St. Benedict's Abbey. P.O. Peramiho, Tanzania, East Africa.
- Bhatt, D., Mitaliya, K., Pandya, N., Baxi, U.S., 2001. Herbal therapy for jaundice. *Advance in Plant Sciences* 14, 123–125.
- Biosafety News, 2002. Environment: Forests Excision could Turn Kenya into Desert. Picasso Productions. Chester House, International Press Center, Koinange Street, P.O. Box 49030, 00100 Nairobi, Kenya.
- Blair, M., Mansingh, A., 2002. Pesticidal potential of tropical plants 111: Acaricidal properties of few crude extracts and their fractions. In: Mansingh, A., Young, E.D., Yee, H.T., Delgoda, R., Robinson, D.E., Morrison, Y.E., Lowe, I.H. (Eds.), *Proceedings of the first international symposium on Utilization of natural products in developing countries: Trends and needs*, held between 10th and 14th July, 2000 at the Natural Products Institute, University of the West Indies, Mona, Kingston 7, Jamaica, West Indies.
- Borges, L.M.F., Ferri, P.H., Silva, W.J., Silva, W.C., Silva, J.G., 2003. In vitro efficacy of extracts of *Melia azedarach* against the tick *Boophilus microplus*. *Medical and Veterinary Entomology* 17, 228–231.
- Botanical South Africa, 2003. Medicinal Uses of Botanicals of South Africa. <http://library.thinkquestfrica.org/TQA00198/treemedical.htm> (27.10.03).
- Bowen, C.R., Hollinger, S.E., 2002. Alternative Crops Web Site, 2nd July, 2002. Illinois State Water Survey, Champaign Illinois. <http://www.sws.uiuc.edu/data/altcrops/> (27.09.03).
- Brechelt, A., 2002. Alternative for natural plant protection in the Dominican Republic. In: Mansingh, A., Young, E.D., Yee, H.T., Delgoda, R., Robinson, D.E., Morrison, Y.E., Lowe, I.H. (Eds.), *Proceedings of the First International Symposium on Utilization of Natural Products in Developing Countries: Trends and Needs*, held between 10th and 14th July, 2000 at the Natural Products Institute, University of the West Indies, Mona, Kingston 7, Jamaica, West Indies.
- Brink, M., 2009. *Grewia trichocarpa* Hochst. ex A.Rich. [Internet] Record from Protabase. In: Brink, M., Achigan-Dako, E.G. (Eds.), *PROTA (Plant Resources of Tropical Africa/Ressources végétales de l'Afrique tropicale)*. Wageningen, Netherlands. <http://database.prota.org/search.htm> (11.11.11).
- Brookes, K.B., Katsoulis, L.C., 2006. Bioactive components of *Rhoicissus tridentata*: a pregnancy-related traditional medicine. *South African Journal of Science* 102, 267–272.

- Buteyo, W.F.G., 2004. Chairman and Co-ordinator, Bungoma Indigenous Trees Conservation Club (BITCC), Bungoma, Kenya.
- Cabral, M.M.O., Garcia, E.S., Rembold, H., De Simone, S.G., Kelecon, A., 1996. Anti-moulting activity in Brazilian *Melia azedarach*. *Memórias Do Instituto Oswaldo Cruz* 91, 117–118.
- Campbell, A., 1994. Community first: Landcare in Australia and London in UK. International Institute for Environment and Development (IIED), Gatekeeper Series. No. 5A42.
- Cassady, J.M., Baird, W.M., Chang, C.J., 1990. Natural products as a source of potential cancer chemotherapeutic and chemopreventive agents. *Journal of Natural Products* 53, 23–41.
- Central Bureau of Statistics, 1996. Kenya Population Census 1989. Analytical Report Vol. 11 April, 1996. Ministry of Planning and National Development, Government of Kenya, Nairobi, Kenya.
- Cevallos-Casals, B.A., Cisneros-Zevallos, L.A., 2002. Bioactive and functional properties of purple sweet potato (*Ipomoea batatas* (L.). *Acta Horticulturae* 583, 195–203. <http://www.actahort.org/books/583/583.22.htm> (21.10.03).
- Chand, N., 1991. Electrical behaviour of sisal fiber. *Indian Journal of Fibre and Textile Research* 16, 287–288.
- Chang, S.G., Chen, C.Y., 1997. Qualitative and quantitative analyses of cannabinoids in *Cannabis sativa*, huo-ma-jen and *Agave sisalana*. *Chinese Pharmaceutical Journal* 49, 77–88.
- Chen, C.C., Chang, S.J., Cheng, L.L., Hou, R.F., 1996. Deterrent effect of the chinaberry extract on oviposition of the diamondback moth, *Plutella xylostella* (L.) (Lep., Yponomeutidae). *Journal of Applied Entomology* 120, 165–169.
- Chifundera, K., 1998. Livestock diseases and the traditional medicine in the Bushi area, Kivu province, Democratic Republic of Congo. *African Study Monographs* 19, 13–33.
- Chih, H.W., Lin, C.C., Tang, K.S., 1996. The hepatoprotective effects of Taiwan folk medicine Ham-hong-chho in rats. *The American Journal of Chinese Medicine* 24, 231–240.
- Ching, F.P., Omogbai, I.K.E., Otokiti, I.O., 2009b. Aqueous stem bark extract of *Stereospermum kunthianum* (Cham, Sandrine Petit) protects against generalized seizures in pentylenetetrazole and electro-convulsive models in rodents. *African Journal of Traditional, Complementary, and Alternative Medicines* 6, 544–548.
- Ching, P.F., Okpo, O.S., Falodun, A., Omogbai, I.K.E., 2009c. Antidiarrhoeal activity of chromatographic fractions of *Stereospermum kunthianum* Cham Sandrine Petit (Bignoniaceae) stem bark. *Tropical Journal of Pharmaceutical Research* 8, 515–519.
- Ching, P.F., Omogbai, I.K.E., Okpo, O.S., Ozolua, I.R., 2009a. Antiinflammatory activity of aqueous extract of *Stereospermum kunthianum* (Cham, Sandrine Petit) stem bark in rats. *Indian Journal of Pharmaceutical Sciences* 71, 106–110.
- Ching, P.F., Omogbai, I.K.E., Ozolua, I.R., Okpo, O.S., 2009d. Analgesic activity of aqueous extract of *Stereospermum kunthianum* (Cham, Sandrine Petit) stem bark. *Acta Poloniae Pharmaceutica* 66, 83–88.
- Chopra, A., Lavin, P., Patwardhan, B., Chitre, D., 2004. A 32-week randomized, placebo-controlled clinical evaluation of RA-11, an ayurvedic drug, on osteoarthritis of the knees. *JCR: Journal of Clinical Rheumatology* 10, 236–245.
- Clevinger, A.J., 2003. Economic and Medicinal Uses. Euphorbiaceae (The Spurge Family). <http://csm.jmu.edu/biology/clevinja/bio486/plantfamilies/AnnSimpson/Euphorbiaceae/euphorbiaceae.htm> (28.11.03).
- Coetzee, C., Jeffthas, E., Reintjen, E., 1999. Indigenous plant genetic resources of South Africa. In: Janick, J. (Ed.), *Perspectives on New Crops and New Uses*. ASHS Press, Alexandria, Virginia, pp. 160–163. [http://www.hort.purdue.edu/Harungana madagascariensis Lam. ex Poir.newcrop/proceedings1999/v4-160.html](http://www.hort.purdue.edu/Harungana%20madagascariensis%20Lam.%20ex%20Poir.newcrop/proceedings1999/v4-160.html) (27.10.03).
- Compaoré, M., Lamien-Meda, A., Mogoşan, C., Lamien, C.E., Kiendrebeogo, M., Voştinaru, O., Vlase, L., Ionescu, C., Nacoulma, O.G., 2011. Antioxidant, diuretic activities and polyphenol content of *Stereospermum kunthianum* Cham. (Bignoniaceae). *Natural Products Research* 25, 1777–1788.
- Cooley, K., Szczerko, O., Perri, D., Mills, E.J., Bernhardt, B., Zhou, Q., Seely, D., 2009. In: Gagnier, J. (Ed.), *Naturopathic Care for Anxiety: A Randomized Controlled Trial* ISRCTN78958974. *PLoS ONE* 4, e6628.
- Corbitt, N.J., Wanyama, E., 2004. Circumcision Rites Among the Babukusu People of Kenya: A Local Christian Response to a Cultural Tradition. Build A Bridge International, Baptist College, Limuru.
- Core, E.L., 1981. Castor bean, *Ricinus communis*. In: McClure, T.A., Lipinsky, E.S. (Eds.), *CRC Handbook of Biosolar Resources. Resource Materials*, vol. 11. CRC Press, Inc., Boca Raton, Florida, pp. 193–197.
- Cremlyn, R., 1978. Pesticides, Preparation and Mode of Action. *Journal of Wiley and Sons*, New York.
- Cunningham, A., Mbenkum, F., 1993. Sustainability of Harvesting *Prunus africana* Bark in Cameroon. A Medicinal Plant in International Trade. People and Plants Working Paper, 2. UNESCO, Paris, France.
- Cunningham, A.B., 1990. Man and Medicines: The Exploitation and Conservation of Traditional Zulu Medicinal Plants. Mitheilungen aus dem Institut für Allgemeine Botanik, Hamburg.
- Cunningham, A.B., 1995. African Medicinal Plants: Setting Priorities at the Interface between Conservation and Primary Healthcare. People and Plants Working Paper 1. <http://www.rbgekew.org.uk/peopleplants/wp/wp1/content.htm> (08.11.03).
- Dagne, E., Alemu, M., Sterner, O., 1993. Flavonoids from *Euclea divinorum*. *Bulletin of the Chemical Society of Ethiopia* 7, 87–92.
- Dalziel, J.M., 1937. *The Useful Plants of West Tropical Africa*. Crown Agents, London, UK.
- Desouter, S., 1991. Pharmacopée humaine et vétérinaire du Rwanda. Musée royal de l'Afrique centrale Tervuren. *Annals of Science Ecology* 22, 254.
- Deweese, A.P., 1995. Social and Economic Incentives for Smallholder Tree Growing. A Case Study from Murang'a District, Central Kenya. Community Forestry Case Study Series. Food and Agriculture Organization of the United Nations, Via delle Terme di Caracalla, 00100 Rome, Italy, <http://www.fao.org/docrep/U8995E/u8995e00.htm#Contents> (28.09.03).
- Ding, Y., Chen, Y.Y., 1989. Steroidal saponins from a cultivated form of *Agave sisalana*. *Phytochemistry* 28, 2787–2792.
- Ding, Y., Tian, R.H., 1993. Two new steroidal saponins from dried fermented residues of leaf juices of *Agave sisalana* forma Dong No. 1. *Chemical and Pharmaceutical Bulletin Tokyo* 41, 557–560.
- Dipeolu, O.O., Mongi, A.O., Punya, D.K., Latif, A.A., Amoo, O.A., Odhiambo, T.R., 1992. Current concepts and approach to control of livestock ticks in Africa. *Discovery and Innovation* 4, 35–44.
- Dipeolu, O.O., Ndungu, J.N., 1991. Acaricidal activity of Kupetaba, a group mixture of natural products, against *Rhipicephalus appendiculatus*. *Veterinary Parasitology* 38, 327–338.
- Dos Santos, G., Miller, R.B., 1997. Wood anatomy of *Jacaranda* (Bignoniaceae): systematic relationships in sections *Monolobos* and *Dilobos* as suggested by twig and stem rays. *IAWA Journal* 18, 369–383.
- Duke, A.J., 1983a. *Euphorbia tirucalli* L. (Euphorbiaceae); Petroleum plant, Aveloz, Milkbrush. In: Duke, A.J. (Ed.), *Handbook of Energy Crops*. Center for New Crops and Plant Products, Purdue University, West Lafayette, India.
- Duke, A.J., 1983. *Handbook of Energy Crops. Acacia nilotica* (L.) Del. (Syn.: *Mimosa nilotica*) commonly called Motse, Egyptian Mimosa, or Thorn. Unpublished work. <http://www.hort.purdue.edu/newcrop/duke.energy/Acacia.nilotica.html> (16.09.04).
- Duke, A.J., 1983. *Handbook of Energy Crops. Sorghum bicolor* (L.) Moench. Unpublished work. <http://www.hort.purdue.edu/newcrop/duke.energy/Sorghum.bicolor.html> (21.10.03).
- Duke, A.J., 2002. A Mini-Course in Medical Botany Syllabus: African Medicinal Plants in World Trade. Module 9: African. <http://www.ars-grin.gov/duke/syllabus/index.html#warn> (06.10.03).
- Duke, A.J., Bogenschütz, J.M., 2003. *Tico Ethnobotanical Dictionary*. Dr. Duke's Phytochemical and Ethnobotanical Databases. USDA, ARS, Genetic Resources. <http://www.ars-grin.gov/duke/dictionary/tico/p.html> (10.11.03).
- Duke, A.J., Wain, K.K., 1981. *Medicinal Plants of the World*. Computer Index with more than 85,000 Entries. 3 Vols.
- Duke-Elder, S., MacFaul, P.A., 1972. Chemical injuries. In: Duke-Elder, S. (Ed.), *System of Ophthalmology. Part 2: Non-mechanical Injuries*, vol. XIV. CV. Mosby Co., St. Louis, Henry Kimpton, London.
- El Tahir, A., Ibrahim, A.M., Satti, G.M.H., Theander, T.G., Kharazmi, A., Khalid, S.A., 1998. The potential antileishmanial activity of some Sudanese medicinal plants. *Phytotherapy Research* 12, 576–579.
- El, T.A., Satti, G.M.H., 1999. Antiplasmodial activity of selected Sudanese medicinal plants with emphasis on *Maytenus senegalensis* (Lam.) Exell. *Journal of Ethnopharmacology* 64, 227–233.
- El-Siddiq, K., Ebert, G., Ludders, P., 1999. Tamarind (*Tamarindus indica* L.): a review on a multipurpose tree with promising future in the Sudan. *Angewandte Botanik* 73, 202–205.
- Erah, O.P., Asonye, C.C., Augustine, O., Okhamafe, O.A., 2003. Response of *Tripinosoma brucei* brucei-induced anaemia to a commercial herbal preparation. *African Journals On-line. African Journal of Biotechnology* 2, <http://www.inasp.org.uk/ajol/journals/ajb/vol2no9abs.html> (29.10.03).
- Etkin, N.L., 1993. Anthropological methods in ethnopharmacology. *Journal of Ethnopharmacology* 38, 93–104.
- Fabry, W., Okemo, P., Aansorg, R., 1996. Activity of East African medicinal plants against *Helicobacter pylori*. *Chemotherapy* 42, 315–317.
- Faninger, A.A., Markovic-Brisk, A., 1960. Occupational dermatosis caused by mouldy sisal. *Srpski arhiv za celokupno lekarstvo* 88, 303.
- Fanshawe, D.B., 1948. Forest products of British Guiana, Part II. Minor Forest Products. *Forestry Bulletin No. 2*. Forestry Department, Georgetown.
- Fanshawe, D.B., 1953. Fish poisons of British Guiana. *Kew Bulletin* 2, 239–240.
- FAO, 1983. Food and Fruit-Bearing Forest Species – Examples from East Africa. *Forestry Paper 44/1*. FAO, Rome, Italy.
- FAO, 1993. Part two: update of recent developments. 2: 4. Processing and utilization. In: *Proceedings of the FAO Expert Consultation on the Role of Forestry in Combating Desertification held in Saltillo, Mexico 24–28 June 1985*, Food and Agriculture Organization (FAO) of the United Nations, Viale delle Terme di Caracalla, 00100 Rome, Italy, <http://www.fao.org/docrep/T0115E/T0115E00.htm#Contents> (09.09.03).
- Fernandes, M.C.E., Oktingati, A., Maghembe, J., 1985. The Chagga Home Gardens: A Multi-storied Agro-forestry Cropping System on Mt. Kilimanjaro, Northern Tanzania. The United Nations University Press, Tokyo, Japan. *Food and Nutrition Bulletin* 7.
- Fox, F.W., Norwood-Young, M.E., 1982. Food from the wild-edible plants of Southern Africa, Delta, Johannesburg, South Africa.
- Franco, L.A., Matiz, G.E., Calle, J., Pinzón, R., Ospina, L.F., 2007. Antiinflammatory activity of extracts and fractions obtained from *Physalis peruviana* L. calyces. *Biomedica* 1, 110–115.
- Freiburghaus, F., Steck, A., Pfander, H., Brun, R., 1998. Bioassay-guided isolation of a diastereoisomer of kolavenol from *Entada abyssinica* active on *Trypanosoma brucei rhodesiense*. *Journal of Ethnopharmacology* 61, 79–183.

- Gachathi, F.N., 1989. Kikuyu botanical dictionary of plant names and uses. Publication supported by GTZ. The print shop, P.O. Box 24576, Nairobi.
- Garneau, F., Gagnon, H., France, J.L., Koumoglou, H. K., Moudachirou, M., Addae-Mensah, I., 1996. Le Chemophytes de Lippia multiflora, Lalaleuca quiquenervia et Clausena anisata Naturels du Togo, Benin et Ghana. Se Colloque Produits Naturels d'Origine Vegetale, Actes du cologne de Saint-Jean sur Richelieu, 18 au 24 Octobre 1995. In: Guy Collin et Francois-Xavier Garneau (Ed.). Laseve University Du Quebec A Chicoutimi, Canada.
- Gathirwa, J.W., Rukunga, G.M., Njagi, E.N.M., Omar, S.A., Mwitari, P.G., Guantai, A.N., Tolo, F.M., Kimani, C.W., Muthaura, C.N., Kirira, P.G., Ndunda, T.N., Amalemba, G., Mungai, G.G.M., Ndiege, O.I., 2008. The *in vitro* anti-plasmodial and *in vivo* anti-malarial efficacy of combinations of some medicinal plants used traditionally for treatment of malaria by the Meru community in Kenya. *Journal of Ethnopharmacology* 115, 223–231.
- Gelfand, M., Mavi, S., Drummond, R.B., Ndemera, B., 1985. The Traditional Medicinal Practitioner in Zimbabwe. Mambo Press, Gweru.
- Getahun, A., 1974. The role of wild plants in the native diet in Ethiopia. *Agro-Ecosystems* 1, 45–56.
- Getahun, A., 1976. Some common medicinal and poisonous plants used in Ethiopian folk medicine. Faculty of Science, Addis Ababa University, Addis Ababa, Ethiopia, p. 63. <http://ip.aaas.org/tekindex.nsf/0/99b535e7618170c485256ae100696f4c/Body/M17OpenElement> (12.10.03).
- Glover, P.E., Stewart, J., Gwynne, M.D., 1966. Maasai and Kipsigis notes on East African plants. Part III – Medicinal uses of plants. *East African Agricultural and Forestry Journal* 32, 200–207.
- Golob, P., Moss, C., Dales, M., Fidge, A., Evans, J., Gudrups, I., 1999. The Use of Spices and Medicinals as Bioactive Protectants for Grains. Natural Resources Institute Chatham Maritime Chatham, Kent, UK. Food and Agriculture Organization of the United Nations, Rome, ISBN 92-5-104294-2, FAO Agricultural Services Bulletin No. 137:M-15.
- Graham, K., Graham, E.A., Towers, G.H.N., 1980. Cercaricidal activity of phenyl heptatriyne and alpha terthienyl naturally occurring compounds in species of Asteraceae compositae. *Canadian Journal of Zoology* 58, 1955–1958.
- Grainge, M., Ahmed, S., 1988. Handbook of Plants with Pest-control Properties. Wiley & Sons, New York.
- Green, E., Obi, C.L., Samie, A., Bessong, O.P., Ndip, N.R., 2011. Characterization of *n*-hexane sub-fraction of *Bridelia micrantha* (Berth) and its antimycobacterium activity. *BMC Complementary and Alternative Medicine* 11, 28–32.
- Grieve, M., 1931. A Modern Herbal. Hafner Press, New York (Reprinted 1974).
- Gruber, W.C., O'Brien, M., 2010. Uterotonic plants and their *bioactive* constituents. *Planta Medica* 77, 207–220.
- Gurumurthy, S.B., Sreenivasa, M.N., 2000. Occurrence and distribution of VAM fungi in the rhizosphere of five tree species grown under agroforestry system. *Environment and Ecology* 18, 500–502.
- Hamburger, M., Marston, A., Hostettmann, K., 1991. Search for new drugs of plant origin. *Advances in Drug Research* 20, 167–215.
- Harjula, R., 1980. Mirau and his practice. A study of the ethnomedicinal repertoire of a Tanzanian herbalist. Institut for cultural research, University of Turku, Finland. Tri-med books limited, London Registered Office, 5 Tudor Cottage, Lovers Walk, Finchley, London, N3 1JH.
- He, W.D., Van Puyvelde, L., Bosselaers, J., De Kimpe, N., Van der Flaas, M., Roymans, A., Mathenge, S., Mudida, F., Chalo Mutiso, P., 2002. Activity of 6-pentadecylsalicylic acid from *Ozoroa insignis* against marine crustaceans. *Pharmaceutical Biology* 40, 74–76.
- He, W.D., Van Puyvelde, L., Bosselaers, J., De Kimpe, N., Van der Flaas, M., Roymans, A., Mathenge, S., Mudida, F., Chalo Mutiso, P., 2000. Bioactive principles from plants of Kenya. IV. Study on the antifungal activity of *Ozoroa insignis* Del. Meded. Fac. Landbouwwet, University of Gent 65, 29–33.
- Heine, B., Heine, I., 1988. Plant concepts and plant use. An ethnobotanical survey of the semi-arid and arid lands of East Africa. Part 1: Plants of the Chamus (Kenya), Band 6, Kolner Beiträge zur Entwicklungsländerforschung/Cologne Development Studies Verlag breitenbach Publishers, Saarbrücken, Fort Lauderdale, Germany, p. 104.
- Heine, B., König, C., 1988. Plant concepts and plant use. An ethnobotanical survey of the semi-arid and arid lands of East Africa. Part 2: Plants of the So (Uganda), Band 7, Kolner Beiträge zur Entwicklungsländerforschung/Cologne Development Studies Verlag breitenbach Publishers, Saarbrücken, Fort Lauderdale, Germany.
- Hempel, P., 2002. Aboriginal traditional knowledge and environmental management. The Science and Environment Bulletin, Environment Canada 32, September/October, 2002 Series.
- Hentgen, A., 1985. Forage trees in India: a chance for the rearing of ruminants. *Fourrages* 101, 105–119.
- HerbWeb, 2000. Global Botanical Exchange. <http://www.herbweb.com> (27.09.03).
- Hines, D.A., Eckman, K., 1993. Indigenous multipurpose trees of Tanzania: uses and economic benefits for people. In: FAO Miscellaneous/93/9 Working Paper, first ed. FAO, Rome/Ottawa, Ontario, Canada, ISBN 0-9697075-0-9, Cultural Survival Canada, Development Services Foundation of Tanzania and Tree Roots Canada <http://www.fao.org/docrep/X5327e/x5327e00.htm> (27.09.03).
- Homer, K.A., Manji, F., Beighton, D., 1990. Inhibition of protease activities of periodontopathic bacteria by extracts of plants used in Kenya as chewing sticks (mswaki). *Archives in Oral Biology* 35, 421–424.
- Homer, K.A., Manji, F., Beighton, D., 1992. Inhibition of peptidase and glycosidase activities of *Porphyromonas gingivalis*, *Bacteroides intermedius* and *Treponema denticola* by plant extracts. *Journal of Clinical Periodontology* 19, 305–310.
- Hussein, G., Miyashima, H., Nakamura, N., Hattori, M., 1999. Inhibitory effects of Sudanese plant extract on HIV-1 replication and HIV-1 protease. *Phytotherapy Research* 13, 31–36.
- Hyde, M.A., Wursten, B.T., Ballings, P., 2011. Flora of Zimbabwe: Cultivated Plants: Species information: *Rhus natalensis*. http://www.zimbabweflora.co.zw/cult/species.php?species_id=136770 (08.12.11).
- Ibrahim, H., Abudurahman, M.E., Shock, M., Ilyas, N., Monday, S., Chordima, W., 1999. Pharmacognostic studies of the stem of *Carrisa edulis*. Online Nigerian Journal of Natural Products and Medicine 3, <http://www.inasp.info/ajol/journals/njnp/vol3abs.html> (20.10.03).
- ICIPE Annual Report, 1998/99. In: Ng'eny-Mengech, A., Sogo, D., Mwanjy, S., Ndung'u, A.C. (Eds.), Sustainable Methods for Integrated Tick Management. The International Centre of Insect Physiology and Ecology (ICIPE), Animal Health Management (Ticks), ICIPE Science Press, Nairobi.
- ICRAF, International Centre for Research in Agroforestry (ICRAF), 1992. A Selection of Useful Trees and Shrubs for Kenya. ICRAF, Nairobi.
- IIRR, 1994. Ethnoveterinary Medicine in Asia: An Information Kit on Traditional Animal Health Care Practices. International Institute of Rural Reconstruction, Silang, Cavite 4118, Philippines.
- Imbabi, E.S., Ibrahim, K.E., Ahmed, B.M., Abulfutuh, I.M., Hulbert, P., 1992. Chemical characterization of tamarind bitter principle, tamarindineal [tamarindineal]. *Fitoterapia* 63, 537–538.
- Irvine, F.R., 1961. Woody Plants of Ghana. With Special Reference to their Uses. Oxford University Press, London.
- Isman, M.B., Matsuura, H., MacKinnon, S., Durst, T., Towers, G.H.N., Arnason, J.T., 1996. Phytochemistry of the Meliaceae. So many terpenoids, so few insecticides. In: Romeo, T.J., Saunders, J.A., Barbosa, P. (Eds.), *Phytochemical Diversity and Redundancy in Ecological Interactions*. Plenum Press, New York.
- ITDG and IIRR, 1996. Ethnoveterinary Medicine in Kenya. A Field Manual of Traditional Animal Health Care Practice. Intermediate Technology Development Group (ITDG) and International Institute of Rural Reconstruction (IIRR), Nairobi, Kenya.
- Izelle, T., 2002. Zulu traditional healers' brew works. MRC News. Science in Africa-Africa's First On-line Science Magazine 29, <http://www.scienceinfrica.co.za/2002/august/brew.htm> (September 2003).
- Jacobson, M., 1990. Glossary of Plant-Derived Insect Deterrents. CRC Press, Inc., Boca Raton, Florida.
- Janet, P., 1986. Trees, plants and a rural community in the southern Sudan. An international Journal of Forestry and Forest Industries. FAO – Food and Agriculture Organization of the United Nations. Unasyula – No. 154 – Agroforestry – pathways, vol. 38, 1986/4. <http://www.fao.org/docrep/50630E/50630e00.htm#Contents> (27.09.03).
- Jawaid, T., Argal, S., Singh, S., 2011. Botanicals and herbs: a traditional approach in treatment of epilepsy. *Journal of Pharmacy Research* 4, 1138–1140.
- Jayasekara, T.K., Belmain, S.R., Stevenson, P.C., Hall, D.R., 2002. *Securidaca longepedunculata* (Fres.) as a control for stored product insect pests. In: Proceedings of the 8th International Working Conference on Stored-Product Protection, York, UK, <http://bru.gmpcr.ksu.edu/proj/iwscpp/04.11.03>.
- Jembere, B., Obeng-Ofori, D., Hassanali, A., Nyamaso, N.N.G., 1995. Products derived from the leaves of *Ocimum kilimandscharicum* (Labiatae) as post-harvest grain protectants against the infestation of three major stored product insect pests. *Bulletin of Entomological Research* 85, 361–367.
- Jeruto, P., Lukhoba, C., Ouma, G., Otieno, D., Mutai, C., 2008. An ethnobotanical study of medicinal plants used by the Nandi people in Kenya. *Journal of Ethnopharmacology* 116, 370–376.
- Johns, T., Faubert, G.M., Kokwaro, J.O., Mahunnah, R.L.A., Kimanani, E.K., 1995. Antigiardial activity of gastrointestinal remedies of the Luo of East Africa. *Journal of Ethnopharmacology* 46, 17–23.
- Johns, T., Kokwaro, J.O., Kimanani, E.K., 1990. Herbal remedies of the Luo of Siaya District, Kenya: establishing quantitative criteria for consensus. *Botanical Botany* 44, 369–381.
- Juliette de Ba Levy, 1991. The Complete Herbal Handbook for Farm and Stable. Faber and Faber Limited, London.
- Juma, C., 1989. Biological Diversity and Innovation: Conserving and Utilizing Genetic Resources in Kenya. ACTS, Nairobi, Kenya.
- Kaaya, P.G., Mwangi, N.E., Malonza, M.M., 1995. Acaricidal activity of *Margaitaria discoidea* (Euphorbiaceae) plant extracts against the ticks *Rhipicephalus appendiculatus* and *Amblyomma variegatum* (Ixodidae). *International Journal of Acarology* 21, 123–129.
- Kambewa, D.M.B., Mftilodze W.M., Hüttner K., Wollny A.B.C., Phoya, D.K.R., 1997. The use of indigenous veterinary remedies in Malawi. In: Mathias-Mundy, E., Rangnekar, V.D., McCorkle, M.C. (Eds.), *Ethnoveterinary Medicine: Alternatives for Livestock Development*. Proceedings of an International Conference held in Pune, India, on November 4–6, 1997. Volume 1: Selected Papers. File 3 of 9: Part 2: Validation of Ethnoveterinary Medicine. BAIF Development Research Foundation, Pune, India.
- Kasonia, K., Ansay, M.N., Basegere, P., Gustin, S., Matamba, M., 1991. Note d'ethnopharmacologie vétérinaire en cas de verminoses, diarrhée, coprostase et météorisme au Kivu et Kibali-Ituri (Zaire). *Tropicicultura* 9, 169–172.
- Kayonga, A., Habiayemye, F.X., 1987. Médecine traditionnelle et plantes médicinales rwandaises. Contribution aux études ethnobotaniques de la flore rwandaise. Préfecture de Gisenyi, Univ. Nat. Rwanda, Centre universitaire de recherche sur la pharmacopée et la médecine traditionnelle, CURPHAMETRA, inédit, Rwanda, p. 121.

- Kebenei, S.J., Ndalut, P.K., Kiprono, C.P., 2004. Larvicidal activity of myricoidine from *Clerodendrum myricoides*. Bulletin of the Chemical Society of Ethiopia 18, 225–227.
- Kennewell, D.P., 1990. Comprehensive Medicinal Chemistry, vol. 1. Pergamon Press, Oxford, London.
- Kenya National Archives (KNA). File No. DC/EN/3/3/2. Annual Abaluyia Land and Law Customs, Nairobi, pp. 1954–1956.
- Killip, E.P., Smith, C.A., 1935. Some American plants used as fish poisons. Catalog Source No. (OCoLC)ocm05794895. Washington: Bureau of Plant Industry and U.S. Department of Agriculture, New York Botanical Garden, New York.
- Kofi-Tsekpo, W.M., Kiyo, D.W., 1998. Newer direction of plant drug research: the need to examine and investigate traditional remedies for pharmacotherapy of animal and zoonotic diseases. African Journal of Health Sciences 5, 12–14.
- Kokwaro, J.O., 1993. Medicinal Plants of East Africa, second ed. East African Literature Bureau, Kampala, Nairobi, Dar Es Salaam.
- Kolar, J., Malbeck, J., 2009. Levels of the antioxidant melatonin in fruits of edible berry species. Planta Medica 75, 9.
- Koumoglou, K.H., Akagana, K., Glitho, A.I., Garneau, F.X., Gagnon, H., Jean, F.I., Moudachirou, M., Addae-Mensah, I., 1996a. Geranial and Neral. Major constituents of *Lippia multiflora* Moldenke leaf oil. Journal of Essential Oil Research 8, 237–240.
- Koumoglou, H.K., Dotse, K., Glotho, A.I., Garneau, F.X., Moudachirou, M., Addae-Mensah, I., 1996b. Essential oils of *Cymbopogon shoenanthus* and *Lippia multiflora* from Togo. Riv. Ital. EPPOS 7, 680–691.
- Kristensen, M., Lykke, M.A., 2003. Informant-based valuation of use and conservation preferences of Savanna trees in Burkina Faso. Economic Botany 57, 203–217.
- Kroschel, J., 1996. Studies on the use of chemicals, botanicals and *Bacillus thuringiensis* in the management of the potato tuber moth in potato stores. Crop Protection 15, 197–203.
- Krueger, R.A., 1988. Focus groups: a practical guide for applied research. In: Agar, M., MacDonald, J. (Eds.), Focus Groups and Ethnography. Human Organization 54, 78–86.
- Kumar, A., 2003. Ayurvedic Medicines: Some Potential Plants for Medicine from India. Energy Plantation Demonstration Project and Biotechnology Center, Department of Botany, University of Rajasthan, Jaipur, 302004 (Rajasthan) India.
- Kunle, O., Okogun, J., Egamana, E., Emojevwe, E., Shok, M., 2003. Antimicrobial activity of various extracts and carvacrol from *Lippia multiflora* leaf extract. Phytomedicine 10, 59–61.
- Ky, K.J.M., 2008. Vitex doniana Sweet. [Internet] Record from Protabase. In: Louppe, D., Oteng-Amoako, A.A., Brink, M. (Eds.), PROTA (Plant Resources of Tropical Africa/Ressources végétales de l'Afrique tropicale), Wageningen, Netherlands. <http://database.prota.org/search.htm> (11.12.11).
- Lan, Y.H., Chang, F.R., Pan, M.J., Wu, C.C., Wu, S.J., Chen, S.L., Wang, S.S., Wu, M.J., Wu, Y.C., 2009. New cytotoxic withanolides from *Physalis peruviana*. Food Chemistry 116, 462–469.
- Lans, C., 2001. Creole remedies. Case Studies of Ethnoveterinary Medicine in Trinidad and Tobago. PhD Thesis, Wageningen University and Research Centre, Wageningen, The Netherlands.
- Lans, C., Brown, G., 1998. Observations on ethnoveterinary medicines in Trinidad and Tobago. Preventive Veterinary Medicine 35, 125–142.
- Le Houérou, H., 2003. FAO Species Description (*Acacia macrothyrsa* Harms.). Grassland and Pasture Crops. <http://www.fao.org/ag/AGP/AGPC/doc/Gbase/DATA/Pf000534.HTM> (12.10.03).
- Lemmens, G.J., Fryer, P.J., 1917. Insecticides, sheep-dip, animal-wash and the like. United States patent, 1,242, 954 and 1,242,955. Patented October 16, 1917.
- Lemordant, D., 1971. Contribution à l'ethnobotanique éthiopienne. Journal d'agriculture tropicale (aterraditionnelle) et de botanique appliqué 18, 142–179.
- Leung, A.Y., Foster, S., 2003. Encyclopedia of Common Natural Ingredients used in Food, Drugs, and Cosmetics, second ed. Wiley-Interscience, The University of Michigan, Ann Arbor, Michigan.
- Lewis, R., 2001. The Institute for Genomic Research (TIGR) Tackles East Coast Fever. Sequencing *Theileria parva* may aid vaccination efforts. The Scientist 15, 16, <http://www.the-scientist.com/yr2001/jan/research.010108.html> (21.06.01).
- Lin, L.F., 1984. Extractions of leaf protein concentrate from sisal (*Agave sisalana*). Journal of Agricultural Research of China 33, 159–168.
- Lindsay, P.J., Kaufman, W.R., 1988. The efficacy of azadirachtin on putative ecdysteroid-sensitive systems in the ixodid tick *Amblyomma americanum* L. Journal of Insect Physiology 34, 439–442.
- Lindsay, R.S., Hepper, F.N., 1978. Medicinal Plants of Marakwet, Kenya. Kew, Royal Botanic Gardens, London.
- Lister, P., 2002. Traditional Indigenous Knowledge. <http://sites.uws.edu.au/vip/listerp/tek.htm> (01.10.03).
- Lukwa, N., Mutambu, S.L., Makaza, N., Molgaard, P., Furu, P., 2001. Perceptions about malaria transmission and control using anti-malaria plants in Mola, Kariba, Zimbabwe. African Journals Online. Nigerian Journal of Natural Products and Medicine 5, <http://www.inasp.info/ajol/journals/njnp/vol5abs.html> (27.10.03).
- Lwande, W., Ndakala, J.A., Hassanali, A., Moreka, L., Nyandat, E., Ndungu, M., Amiani, H., Gitu, P.M., Malonza, M.M., Punyua, K.D., 1999. Gynandropsis gynandra (L) Brig. essential oil and its constituents as tick (*Rhipicephalus appendiculatus*) repellents. Phytochemistry 50, 401–405.
- Mahgoub, S.M., Ahmed, S.M., 1996. *Ricinus communis* seed extract as protectants of wheat grains against the rice weevil, *Sitophilus oryzae* L. Annals of Agricultural Science, Cairo 41, 483–491.
- Makila, F.E., 1978. An Outline of History of the Babukusu. Kenya Literature Bureau, Nairobi, Kenya.
- Malerich, P.J., Trauner, D., 2003. Biomimetic synthesis of (±)-Pinnatal and (±)-Stereokunthal A. Journal of American Chemical Society 125, 9554–9555.
- Malonza, M.M., Dipeolu, O.O., Amoo, A.O., Hassan, S.M., 1992. Laboratory and field observations on anti-tick properties of the plant *Gynandropsis gynandra* (L) Brig. Veterinary Parasitology 42, 123–136.
- Mann, E., 2001. Medicinal trees of Bukusuland. In: Mann, E. (Ed.), Council for Human Ecology, Kenya (CHECK), first ed. Annex Import, Nairobi.
- Mansingh, A., Williams, L.A.D., 1998. Pesticidal potential of tropical plants – II. Acaricidal activity of crude extracts of several Jamaican plants. Insect Science and Applications 18, 149–155.
- Mansingh, A., Williams, L.A.D., 2002. Pesticidal potential of tropical plants – VIII. Acute toxicity and inhibition of oogenesis in the ticks *Boophilus microplus* Canest. and *Amblyomma cajennense* Fab. by the extracts of *Azadirachta indica* A. Juss and *Artocarpus altilis* Park. In: Mansingh, A., Young, E.D., Yee, H.T., Delgoda, R., Robinson, D.E., Morrison, Y.E., Lowe, I.H. (Eds.), Proceedings of the First International Symposium on Utilization of Natural Products in Developing Countries: Trends and Needs, held between 10th and 14th July, 2000 at the Natural Products Institute, University of the West Indies, Mona, Kingston 7, Jamaica, West Indies.
- Marin, R., 1999. Collection of Medicinal Plant Samples from Africa and Elsewhere. World Botanical Associates, <http://www.worldbotanical.com/WBAMED2.HTM> (01.10.03).
- Martin, J.G., 1996. Ethnobotany, A 'people and Plants' Conservation Manual. Chapman & Hall, London, New York.
- Martin, M., Mathias, E., McCorkle, M.C., 2001. Ethnoveterinary Medicine: An Annotated Bibliography of community Animal Healthcare, second ed. ITDG Publishing, London.
- Martina, M.B., 2001. The role of indigenous trees for the conservation of biocultural diversity in traditional agroforestry land use systems: the Bungoma case study: *in situ* conservation of indigenous tree species. Agroforestry Systems 52, 119–132.
- Martina, M.B., 1998. Indigenous Trees in Bukusuland. A Selection of Useful Trees and Shrubs for Social Forestry in Bungoma. Executive Printing Works Limited, Nairobi.
- Masinde, P.S., 1996. Medicinal plants of the Marachi people of Kenya. The biodiversity of African plants. In: van der Maesen, J.G., van der Burgt, X.M., van Medenbach de Rooy, J.M. (Eds.), Proceedings of the 14th AETFAT Congress, Wageningen, The Netherlands, 22–27 August 1994. Kluwer Academic Publishers, The Netherlands.
- Mathias-Mundy, E., McCorkle, M.C., 1989. Ethnoveterinary Medicine: An annotated bibliography. Bibliographies in Technology and Social Change, No. 6, Technology and Social Change Programme, IOWA State University, Ames, IOWA 50011 USA.
- Matzigkeit, U., 1990. Natural Veterinary Medicine. Ectoparasites in the Tropics. Josef Margraf Verlag, Weikersheim.
- Maundu, P.M., Ngugi, G.W., Kabuye, C.H.S., 1999. Traditional Food Plants of Kenya, first ed. Kenya Resource Centre for Indigenous Knowledge (KENRIK), National Museums of Kenya, English Press Limited, Nairobi, ISBN 9966-9861-4-6.
- Maundu, P.M., Berger, D., Ole Saitabau, J., Nasieku, C., Kipelian, J., Mathenge, M., Morimoto, S.G., Hoft, Y., 2001. Ethnobotany of the Loita Maasai: Towards Community Management of the forest of the Lost Child-Experiences from the Loita Ethnobotany Project. People and Plants Working Paper 8. UNESCO, Paris, France. <http://www.rbgekew.org.uk/peopleplants/wp/wp8/index.html> (28.09.03).
- McCorkle, M.C., Rangnekar, V.D., Mathias-Mundy, E., 1997. Introduction: whence and whither ER&D? In: Mathias, E., Rangnekar, V.D., McCorkle, M.C., Martin, M. (Eds.), Ethnoveterinary Medicine: Alternatives for Livestock Development. Proceedings of an International Conference held in Pune, India, November 4–6, 1997. Volume 1: Selected Papers. File 2 of 9: Part 4: Introduction and Part 1: Applied Studies of Ethnoveterinary Systems. BAIF Development Research Foundation, Pune, India.
- McIntyre, A., 2002. Ashwagandha: Winter Cherry – *Withania somnifera*. <http://www.positivehealth.com/permit/Articles/Regular/mcintyre77.htm> (21.10.03).
- Michael, H.M., 2003. Medicinal Uses of Succulents. <http://www.succulents.co.za/Contact.shtml>. November 8, 2003, 11:08AM East Africa Time.
- Minja, M.M.J., 1989. Collection of Tanzanian medicinal plants for biological activity studies. In: Proceedings of the 7th Tanzania Veterinary Association Scientific Conference, vol. 7, Tanzania Veterinary Association, pp. 67–78.
- Minja, M.M.J., 1994. Medicinal Plants used in Promotion of Animal Health in Tanzania. Animal Diseases Research Institute, Department of Pharmacology and Toxicology, Dar Es Salaam, Tanzania.
- Mirjalili, M.H., Moyano, E., Bonfill, M., Cusido, R.M., Palazón, J., 2009. Steroidal lactones from *Withania somnifera*, an ancient plant for novel medicine. Molecules 14, 2373–2393.
- Mitchell, S.M., Ahmad, M.H., 2006. A review of medicinal plant research at the University of the West Indies, Jamaica, 1948–2001. West Indian Medical Journal 55, 243–269.
- Mizrahi, Y., Nerd, A., Sitrit, Y., 2002. New fruits for arid climates. In: Janick, J., Whipkey, A. (Eds.), Trends in New Crops and New Uses. ASHS Press, Alexandria, Virginia, pp. 378–384.
- Morgan, D.L., 1988. Focus groups as qualitative research. In: Agar, M., MacDonald, J. (Eds.), Focus Groups and Ethnography. Human Organization 54, 78–86.
- Morton, J., 1987. Cape gooseberry. In: Morton, J.F., Miami, F.L. (Eds.), Fruits of Warm Climates. <http://www.hort.purdue.edu/newcrop/morton/cape-gooseberry.html> (03.11.03).
- Morton, J.F., 1977. Poisonous and injurious higher plants and fungi. In: Tedeschi, C.G. (Ed.), Forensic Medicine 3, 1456. Philadelphia, Saunders.

- Morton, J.F., 1981. Atlas of Medicinal Plants of Middle America. Bahamas to Yucatan. C.C. Thomas, Springfield, Illinois.
- MTDP, 2003. Molecular Targets Development Program (MTDP). Center for Cancer Research at the National Cancer Institute, <http://home.ncicrf.gov/mtdp/index.html>. October 20, 2003. 3:05PM East Africa Time.
- Muhammad, I., Mossa, J.S., Mirza, H.H., El-Feraly, F.S., 1999. A new modified 6,7-secolabdane diterpenoid from *Clusia richardiana* L. *Phytochemistry* 50, 1225–1227.
- Mulholland, D.A., Monkhe, T.V., Coombes, P.H., Rajab, M., 1998. Limonoids from *Turraea holstii* and *Turraea floribunda*. *Phytochemistry* 49, 2585–2590.
- Mulla, M.S., Su, T., 1999. Activity of biological effects of neem products against arthropods of medical and veterinary importance. *Journal of American Mosquito Control Association* 15, 133–152.
- Mutangah, J.G., 1996. Forest quality indicators with reference to Kakamega forest, Kenya. Plant Conservation Programme, National Museums of Kenya. <http://www.earthwatch.org/europe/limbe/upf.html#top> (18.10.03).
- Mwangi, E.N., 1996. The role of women in animal husbandry: the Kenyan perspective. In: Zessin, H.K. (Ed.), *Livestock Production and Diseases in the Tropics: Livestock Production and Human Welfare. Proceedings of the VIII International Conference of Institutions of Tropical Veterinary Medicine held from the 25th to 29th September, 1995, Berlin, Germany. Volume II. Deutsche stiftung für Internationale Entwicklung. Zentralstelle für Ernährung und Landwirtschaft, Feldafing, Germany.*
- Mwangi, J., Thoithi, G.N., Addae-Mensah, I., Achenbach, H., Lwande, W., Hassanali, A., 1998. Aromatic plants of Kenya III: volatile and some non-volatile constituents of *Croton sylvaticus* Hochst. East and Central African Journal of Pharmaceutical Sciences 1, 41–43.
- Mwangi, J.W., Addae-Mensah, I., Lwande, L.W., Munavu, R.M., 1991b. Essential oils of two *Lippia ukambensis* (Vatke) chemophytes and *Lippia somalensis* (Vatke) in Kenya. *Journal of Essential oil Research* 3, 413–417.
- Mwangi, J.W., Addae-Mensah, I., Munavu, R.M., Lwande, L., 1995. The potential for commercialization of three African *Lippia* species as sources of Essential oil for perfumery and medicinal purposes. In: *Volatilisation de la Biomasse Végétale Par les Produits Naturels*, (Actes du Colloque de Chicoutimi, 22 au 25 Aout 1993. Université du Québec à Chicoutimi, Canada) Published by Centre de Recherches Pour le Développement International (CRDI/IDRC), pp. 205–216.
- Mwangi, J.W., Addae-Mensah, I., Muriuki, G., Lwande, W., Munavu, R.M., 1991a. Essential oils of Kenya *Lippia* species III. Flavour and Fragrance Journal 6, 221–224.
- Mwangi, J.W., Muriuki, G., Addae-Mensah, I., Munavu, R.M., Lwande, W., Cravero, A.A., Alencar, J.W., 1989. Essential oils of *Lippia wilmsii*. *Ev. Latinoamer Ouim.* 20, 143–144.
- Mwangi, J., Njonge, E.W., Addae-Mensah, I., Munavu, R.M., Lwande, W., 1994. Antimicrobial activity of the essential from *Lippia* species in Kenya. *Discovery and Innovation* 6, 58–60.
- Mweseli, M., 2004. Art, Wit and Wisdom of African Proverbs – Selected 'Bukusu' Proverbs from Kenya, their translation into Swahili and English and their Meanings. In: *International Society for Oral Literature in Africa (ISOLA) Fifth Conference of ISOLA, 15–17 July 2004, University of the Gambia, Banjul.*
- Nangendo, S.M., 1994. Daughters of the Clay, Women of the Farm: Women, Agricultural Economic Development, and Ceramic Production in Bungoma District, Western Province, Kenya. Unpl. Ph.D. Dissertation, Department of Anthropology, Bryn Mawr College, Pennsylvania, USA.
- Nardo, E.A.B., Costa, A.S., Lourencao, L.A., 1997. *Melia azedarach* extract as an antifeedant to *Bemisia tabaci* (Homoptera: Aleyrodidae). *Florida Entomologist* 80, 92–94.
- Nasimiyu, R., 1985. The participation of women in Bukusu economy: the situation as at the end of nineteenth century. In: Wandibba, S. (Ed.), *History and Culture in Western Kenya: The People of Bungoma District Through Time*. Gideon S. Were Press, Nairobi, pp. 51–64.
- Ndumu, P.A., George, J.B.D., Choudhury, M.K., 1999. Toxicity of neem seed oil (*Azadiracta indica*) against the larvae of *Amblyomma variegatum* a three-host tick in cattle. *Phytotherapy Research* 13, 532–534.
- Ndung'u, M.W., Chhabra, C.S.W., Lwande, W., 1995. Cleome hirta essential oil as livestock tick (*Rhipicephalus appendiculatus*) and maize weevil (*Sitophilus zeamais*) repellent. *Fitoterapia* 70, 514–516.
- Ndung'u, M.W., Lwande, W., Hassanali, A., Moreka, L., Chhabra, C.S., 1999. Cleome monophylla essential oil and its constituents as tick (*Rhipicephalus appendiculatus*) and maize weevil (*Sitophilus zeamais*) repellents. *Entomologia Experimentalis et Applicata* 76, 217–222.
- Neupane, F.P., 1992. Insect pests associated with some fuelwood and multipurpose tree species in Nepal. *Journal of Tropical Forest Science* 5, 1–7.
- Neuwinger, H.D., 2000. African Traditional Medicine: A Dictionary of Plant Use and Applications. Medpharm Scientific, Stuttgart.
- New Agriculturist, 2003. Reporting Agriculture for the 21st Century. From maths matrix game to management tools. www.new-agri.co.uk/00-3/develop/dev04.html (10.10.03).
- Niang, A., 1987. Contribution à l'étude de la Pharmacopée traditionnelle mauritanienne. Thèse pour le doctorat en médecine vétérinaire. Ecole Nationale de Médecine vétérinaire, SIDI, THABET, Tunisie, p. 156.
- Nobel, P.S., Berry, W.L., 1985. Element responses of *Agave*. *American Journal of Botany* 72, 686–694.
- Nyazema, Z.N., 2002. Evaluation of phytotherapy (Herbalism). Department of Clinical Pharmacology, Medical School. P.O. Box A 178, Avondale, Harare, Zimbabwe. <http://groups.yahoo.com/group/Phytomedica/message/1399?> (08.11.03).
- Oakley, A., 1981. Interviewing women: a contradiction in terms. In: De Rocheleau, D. (Ed.), *Participatory Research and the Race to Save the Planet: Questions, Critique and Lessons from the Field. Agriculture and Human Values* 11, 4–25.
- Odda, J., Kristensen, S., Kabasa, J., Waako, P., 2008. Larvicidal activity of *Combretum collinum* Fresen against *Aedes aegypti*. *Journal of Vector Borne Diseases* 45, 321–324.
- Ogundaini, A., 1999. Antimicrobial agents from some Nigerian plants. *African Journals On-line. Nigerian Journal of Natural Products and Medicine* 3, <http://www.inasp.info/ajol/journals/njnp/vol3abs.html> (20.10.03).
- Okwemba, A., 2002. Local importance herb to flavour yoghurt. An AWC Feature. *Horizon Magazine*. <http://www.nationaudio.com/News/DailyNation/Supplements/horizon/24102002/story2.htm> (17.10.02).
- Olusola, L., Zebulon, S.C., Okoye, U.F., 1997. Effect of *Vitex doniana* stem bark on blood pressure. *Nigerian Journal of Natural Products and Medicine*, <http://www.inasp.info/ajol/journals/njnp/vol1abs.html> (08.11.03).
- Onegi, B., Kraft, C., Kohler, I., Freund, M., Jenett-Siems, K., Siems, K., Beyer, G., Melzig, M.F., Bienze, U., Eich, E., 2002. Antiplasmodial activity of nephthoquinones and one anthraquinone from *Stereospermum kunthianum*. *Phytochemistry* 60, 39–44.
- Onge, St. J., 2002. Fish-poison Use in the Americas. <http://www.survival.com/fish.htm> (10.11.03).
- Oni, P.I., 2001. State of Forest Genetic Resources in the dry north of Nigeria. Sub-Regional Workshop FAO/IPGRI/ICRAF on the conservation, management, sustainable utilization and enhancement of forest genetic resources in Sahelian and North-Sudan Africa (Ouagadougou, Burkina Faso, 22–24 September, 1998). Forest Genetic Resources Working Papers, Working Paper FGR/16E. Forestry Department, FAO, Rome, Italy.
- Orwa, C., Mutua, A., Kindt, R., Jamnadass, R., Simons, A., 2009. Agroforestry Database: A Tree Reference and Selection Guide Version 4.0. <http://www.worldagroforestry.org/af/treedb/> (11.12.11).
- Palgrave, K.C., 1988. Trees of Southern Africa, Revised Edition. C. Struik Publishers, Cape Town/Johannesburg, South Africa.
- Pande, B.C., 1999. Zeetress – a promising anti-stress remedy and immunomodulator. A review. In: Mathias, E., Rangnekar, V.D., McCorkle, M.C. (Eds.), *Ethnoveterinary Medicine: Alternatives for Livestock Development. Proceedings of an International Conference held in Pune, India, on November 4–6, 1997. Volume 1: Selected Papers. BAIF Development Research Foundation, Pune, India.*
- Pardo, J.M., Fontanilla, M.R., Ospina, L.F., Espinosa, L., 2008. Determining the pharmacological activity of *Physalis peruviana* fruit juice on rabbit eyes and fibroblast primary cultures. *Investigative Ophthalmology and Visual Science* 7, 3074–3079.
- Perry, L.M., 1980. Medicinal Plants of East and Southeast Asia. MIT Press, Cambridge.
- Persson, J., 1986. Trees, plants and a rural community in the southern Sudan. *International Journal of Forestry and Forest Industries* 38, Unasylva, FAO <http://www.fao.org/docrep/50630E/50630e00.htm#Contents> (10.11.03).
- Petit, S., 2003. FAO Species Description (*Stereospermum kunthianum* Cham.). Grassland and pasture crops. <http://www.fao.org/ag/AGP/AGPC/doc/Cbase/DATA/Pf000534.HTM> (10.10.03).
- Philippe and Sura Faucon, 2003. Desert Tropics. <http://www.desert-tropics.com> (24.09.03).
- Phillipson, P., 1995. A Leaf from the Herbarium. Notes from the Selmar Schonland Herbarium, Grahamstown, South Africa. Issue No. 2, September 1995.
- Pinto, M.D.S., Ranilla, L.G., Apostolidis, E., Lajolo, F.M., Genovesi, M.I., Shetty, K., 2009. Evaluation of antihyperglycemia and antihypertension potential of native Peruvian fruits using *in vitro* models. *Journal of Medicinal Food* 12, 278–291.
- Pires, M.D.F.C., Purchio, A., 1991. Mycological evaluation of filtrated juice of *Agave sisalana* Perrine's leaf. *Revista De Microbiologia* 22, 272–275.
- Politz, J., Lekeleley, J., 1988. The knowledge of the Samburu on animal diseases and their traditional methods of treatment. Unpublished research work.
- Polygenis-Bigendako, M.J., 1990. Recherches ethnopharmacognosiques sur les plantes utilisées en médecine traditionnelle au Burundi occidental. Thèse présentée en vue de l'obtention du grade de Docteur en sciences, année acad. 1989–1990. Université libre de Bruxelles, Faculty of Science Laboratory. de Botanique systématique et de Phytosociologie.
- Port, R., 2000. Some Local History of the Bungoma Area. Kenya Friendship. The Committee to Support Links between Bloomington, Indiana and Bungoma, Western Province, Kenya. <http://www.cs.indiana.edu/~port/kenyafriends/index.html> (19.05.04).
- Puckhaber, L.S., Stipanovic, R.D., Bost, G.A., 2002. Analyses for flavonoid aglycones in fresh and preserved *Hibiscus* flowers. In: Janick, J., Whipkey, A. (Eds.), *Trends in New Crops and New Uses. ASHS Press, Alexandria, Virginia*, pp. 556–563. <http://www.hort.purdue.edu/newcrop/ncnu02/v5-556.html> (21.10.03).
- Puyvelde (van), L., Geysen, D., Ayobangira, F.X., Hakizamungu, E., Nshimiyanima, A., Kalisa, A., 1985. Screening of medicinal plants of Rwanda for acaricidal activity. *Journal of Ethnopharmacology* 13, 209–215.
- Rahmatullah, M., Samarrai, W., Jahan, R., Rahman, S., Sharmin, N., Miajee, U.E.M.U.Z., Chowdhury, H.M., Bari, S., Jamal, F., Bashir, A.M.B.A., Azad, K.A., Ahsan, S., 2010. An ethnomedicinal, pharmacological and phytochemical review of some Bignoniaceae family plants and a description of Bignoniaceae plants in folk medicinal uses in Bangladesh. *Advances in Natural and Applied Sciences* 4, 236–253.
- Rajab, S., Bentley, M.D., Hassanali, A., Chapya, A., 1988. A new limonoid from *Turraea holstii*. *Phytochemistry* 27, 2353.
- Rajab, S.M., Mulholland, A.D., Monkhe, T.V., 1998. Triterpenoids from *Turraea holstii*. *Phytochemistry* 49, 2585.
- Ramos-Escudero, F., Mu Oz, A.M., Alvarado-Ort, Z.C., Alvarado, N., Y Ez, J.A., 2011. Purple corn (*Zea mays* L.) phenolic compounds profile and its assessment as an

- agent against oxidative stress in isolated mouse organs. *Journal of medicinal food* (Epub. ahead of print) <http://www.ncbi.nlm.nih.gov/pubmed/22082063> (12.12.11).
- Rao, Y.S., Mathew, K.M., Potty, S.N., 1999. Tamarind (*Tamarindus indica* L.) research – a review. *Indian Journal of Arecanut, Spices and Medicinal Plants*, 127–145.
- Regassaa, A., 2000. The use of herbal preparations for tick control in western Ethiopia. *Journal of South Africa Veterinary Association* 71, 240–243.
- Roach, R.G., 2003. People Profile: The Luhya of Kenya. Baptist Mission of Kenya: Orville Boyd Jenkins. <http://www.geocities.com/orvillejenkins/profiles/Luhya.html> (15.05.04).
- Rojas, J., Ochoa, V., Ocampo, S., Muñoz, J., 2006. Screening for antimicrobial activity of ten medicinal plants used in Colombian folkloric medicine: a possible alternative in the treatment of non-nosocomial infections. *BMC Complementary and Alternative Medicine* 6, 2.
- RSCU, 1992. A Selection of Useful Trees and Shrubs for Tanzania. Regional Soil Conservation Unit (RSCU), Draft, Nairobi, Kenya.
- Rukangira, E., 2001. The African Herbal Industry: Constraints and Challenges. *Erboristeria Domani*. http://www.conserveafrica.org/herbal_industry.rtf (04.11.03).
- Rukangira, E., 2003. Overview on the Importance and the Status of Medicinal Plants and Traditional Medicine in Africa. Sustainable Development International. <http://www.sustdev.org/Features/04.09.01.shtml> (27.10.03).
- Rulangaranga, Z.K., 1989. Some Important Indigenous Medicinal and Aromatic Plants in the Wild Flora of Tanzania Mainland. Tropical Forestry Action Plan, Working Paper 24. Tanzania Ministry of Lands, Natural Resources and Tourism, Dar es Salaam, Tanzania.
- Sahney, S., Benton, M.J., 2008. Recovery from the most profound mass extinction of all time. *Proceedings of the Royal Society B: Biological Sciences* 275, 759–765.
- Sahney, S., Benton, M.J., Ferry, P.A., 2010. Links between global taxonomic diversity, ecological diversity and the expansion of vertebrates on land. *Biology Letters* 6, 544–547.
- Saka, J.D.K., Msonthi, J.D., 1994. Nutritional value of indigenous wild trees in Malawi. *Forest Ecology Management* 64, 245–248.
- Samuelsson, G., Faraha, M.H., Claeson, P., Hagos, M., Thulin, M., Hedberg, O., Warfa, A.M., Hassan, A.O., Elmi, A.H., Abdurahman, A.D., Elmi, A.S., Abdie, Y.A., Alin, M.H., 1992. Inventory of plants used in traditional medicine in Somalia. II. Plants of the families Combretaceae – Labiatae. *Journal of Ethnopharmacology* 37, 47–70.
- Sanon, S., Azas, N., Gasquet, M., Ollivier, E., Mahiou, V., Barro, N., Cuzin-Ouattara, N., Traore, A.S., Esposito, F., Balansard, G., Timon-David, P., 2003. Antiplasmodial activity of alkaloid extracts from *Pavetta crassipes* (K. Schum) and *Acanthospermum hispidum* (DC), two plants used in traditional medicine in Burkina Faso. *Parasitology Research* 90, 314–317.
- Scartezzini, P., Speroni, E., 2000. Review on some plants of Indian traditional medicine with antioxidant activity. *Journal of Ethnopharmacology* 71, 23–43.
- Schippmann, U., 2001. Medicinal plants significant trade. CITES Projekt S-109, Plants Committee Document PC9 9.1.3 (rev.). BFN Scripten – 39, BFN-German Federal Agency for Nature Conservation, pp. 51–58.
- Schmidt, G.H., Rembold, H., Ahmed, A.A.I., Breuer, M., 1998. Effect of *Melia azedarach* fruit extract on juvenile hormone titer and protein content in the hemolymph of two species of noctuid lepidopteran larvae (Insecta: Lepidoptera: Noctuidae). *Phytoparasitica* 26, 283–292.
- Schmidt, J.R., 2003. Agavaceae (Agave or Century Plant family). Plant Families Index, The Botanical Dermatology Database (BoDD), Index to Botanical Dermatology Reviews. <http://BoDD.cf.ac.uk/BoDDHomePage.html> (09.10.03).
- Schmidt, J.R., 2003. Euphorbiaceae (Spurge family). Plant Families Index, The Botanical Dermatology Database (BoDD), Index to Botanical Dermatology Reviews. <http://BoDD.cf.ac.uk/BoDDHomePage.html> (29.09.03).
- Schmidt, J.R., 2003. Guttiferae (St John's Wort family). Plant Families Index, The Botanical Dermatology Database (BoDD), Index to Botanical Dermatology Reviews. <http://BoDD.cf.ac.uk/BoDDHomePage.html> (29.10.03).
- Schmidt, J.R., 2003. Leguminosae (Pea family). Plant Families Index, The Botanical Dermatology Database (BoDD), Index to Botanical Dermatology Reviews. <http://BoDD.cf.ac.uk/BoDDHomePage.html> (18.10.03).
- Schwartz, L., Tulipan, L., Birmingham, D.J., 1957. Occupational Diseases of the Skin, third ed. Henry Kimpton, London.
- Seo, Y., 2002. A new bioactive triterpenoid saponin from *Pittosporum viridiflorum* from the Madagascar Rainforest. *Journal of Natural Products* 65, 65–68.
- Sika, F.K.N., 1996. Behavioural responses of *Rhipicephalus appendiculatus* Neumann 1901 and *R. evertsi* Neumann 1889 (Acari: Ixodidae) to host and non-host semiochemicals. PhD Thesis, Zoology Department, Kenyatta University, Nairobi, Kenya.
- Simon, F., 1998. Bioprospecting or biopiracy? *Drug Discovery Today* 3, 399–402.
- Singh, P.R., 2002. Strategies for the management of crop pests with neem (*Azadirachta indica* A. Juss.) and other plant-derived insecticides. In: Mansingh, A., Young, E.D., Yee, H.T., Delgoda, R., Robinson, D.E., Morrison, Y.E., Lowe, I.H. (Eds.), *Proceedings of the First International Symposium on Utilization of Natural Products in Developing Countries: Trends and Needs*, held between 10th and 14th July, 2000 at the Natural Products Institute, University of the West Indies, Mona, Kingston 7, Jamaica, West Indies.
- Singh, S., Kumar, S., 1998. *Withania somnifera*. The Indian Ginseng Ashwagandha. Lucknow. ISBN: 81-86943-91-9. <http://www.vedamsbooks.com/no15185.htm> (21.10.03).
- Smith, F.P., 1969. Chinese Materia Medica. Vegetable kingdom, second ed. (revised by Hei, P.D.). Ku Ting Book House, Taipei, Taiwan.
- Ssegawa, P., Kasenene, M.J., 2007. Medicinal plant diversity and uses in Sango bay area, Southern Uganda. *Journal of Ethnopharmacology* 113, 521–540.
- Staner, P. and Boutique, R. 1937. Matériaux pour l'étude des plantes médicinales indigènes du Congo Belge. *Mém. Institut royal colonial belge, Section des Sc. naturelles et médicales, Collection in-8°, fasc. 6 et dernier*, 228 pp.
- Stedman, T.L., Wilcox, M.D., Follett, Co., 1942. *Stedman's Shorter Medical Dictionary*. <http://www.botanical.com/botanical/mgmh/n/nuxvom08.html> (22.08.04).
- Stockbauer, B., 2003. Tree list for environmental nurseries. African Blackwood Conservation Project. <http://www.blackwoodconservation.org/5-year-program.tree.list.html> (18.10.03).
- Stoll, G., 1988. *Natural Plant Protection in the Tropics*. AGRECOL, Margraf Publishers Scientific Books, Weikersheim, Germany.
- Sutton, J.A., Orr, B.D., 1991. The use of the school essay as an RRA technique: A case study from Bond County, Liberia. *RRA Notes. Participatory Methods for Learning and Analysis*. No. 14. IIED Sustainable Agriculture Program.
- Tadesse, D., 1994. Traditional use of some medicinal plants in Ethiopia. In: Seyani, J.H., Chicuni, A.C. (Eds.), *Proceedings of the 13th Plenary Meeting of AETFAT, Zomba, Malawi, (April 2–11, 1991)*, vol. 1. National herbarium and Botanic Gardens of Malawi, Zomba, pp. 273–293.
- Taylor, F.W., 1986. The potential for utilization of indigenous plants in Botswana. In: Wickens, G.E., Goodin, J.R., Field, D.V. (Eds.), *Plants for Arid Lands*. George Allen and Unwin, London, pp. 231–242.
- Taylor, L., 2002. *Herbal Secrets of the Rainforest*, second ed. In: Taylor, L. (Ed.), *Tropical Plant Database. Database File for Picão Preto (Bidens pilosa)*. Sage Press Inc., P.O. Box 80064, Austin, TX 78708-0064. <http://www.rain-tree.com/picaopreto.htm> (01.11.03).
- Thapa, J.M., Pathak, N.L., 2002. The medicinal and health food plants of Nepal. In: Mansingh, A., Young, E.D., Yee, H.T., Delgoda, R., Robinson, D.E., Morrison, Y.E., Lowe, I.H. (Eds.), *Proceedings of the First International Symposium on Utilization of Natural Products in Developing Countries: Trends and Needs*, held between 10th and 14th July, 2000 at the Natural Products Institute, University of the West Indies, Mona, Kingston 7, Jamaica, West Indies.
- The World Bank Group, 2003. Indigenous knowledge Program for Development. Knowledge Pack: Health. <http://www.worldbank.org/afr/ik/ikpacks/health.htm>, as retrieved October 29th, 2003.
- Thijssen, H.J.C., 2008. *Vitex fischeri* Gürke. [Internet] Record from Protabase, in: Louppe, D., Oteng-Amoako, A.A., Brink, M. (Eds.), *PROTA (Plant Resources of Tropical Africa/Ressources végétales de l'Afrique tropicale)*, Wageningen, Netherlands. <http://database.prota.org/search.htm> (11.11.11).
- Timberlake, J.R., 1987. Ethnobotany of the Pokot of Northern Kenya. Unpublished report. East African Herbarium, Nairobi, Kenya and Center for Economic Botany, Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AE, UK, p. 106.
- Torto, B., Hassanali, A., 1997. Progress in the search for anti-arthropod botanicals. *Recent Research Development in Phytochemistry* 1, 475–488.
- Toyang, J.N., Nuwayyakpa, M., Ndi, C., Django, S., Kinyuy, C.W., 1995. Ethnoveterinary medicine practices in the Northwest Province of Cameroon. *IK Monitor* 3, <http://www.nuffic.nl/ciran/ikdm/3-3/articles/toyang.html> (20.10.03).
- Trade Winds Fruit, 2003. <http://www.tradewindsfruit.com/custservice.htm> (26.10.03).
- Tredgold, M.H., 1990. *Food Plants of Zimbabwe*. Mambo Press, Harare, Zimbabwe (Reprinted 1990).
- UNEP World Conservation Monitoring Centre, 2003. *Traditional Lifestyles and Biodiversity Use. Regional Report (UNEP/CBD/WG8/3/INF/3): AFRICA*. Composite Report on the Status and Trends Regarding the Knowledge, Innovations and Practices of Indigenous and Local Communities Relevant to the Conservation and Sustainable Use of Biodiversity.
- USDA, ARS, National Genetic Resources Program, 2004. *Phytochemical and Ethnobotanical Databases* [Online Database]. National Germplasm Resources Laboratory, Beltsville, Maryland, USA, <http://www.arsgrin.gov:8080/ngpspub/xsql/duke/pl.act.xsql?taxon=2160> (04.03.04).
- UWVG, 2003. Uganda Wildlife Medicinal Garden (UWVG). A list of some of over 500 herbs in Uganda Wildlife Medicinal Garden, with information about diseases they cure. http://www.uweczo.org/Medicinal_plants.tree.tags.pdf (09.11.03).
- Valladares, G., Defago, M.T., Palacios, S., Carpinella, M.C., 1997. Laboratory evaluation of *Melia azedarach* (Meliaceae) extracts against the elm leaf beetle (Coleoptera: Chrysomelidae). *Journal of Economic Entomology* 90, 747–750.
- van Anel, T., 2000. The diverse uses of fish-poison plants in northwest Guyana. *Economic Botany* 54, 500–512.
- Van Der Hoof, C.S., Hoekstra, A., Winter, A., De Smet, P.A., Stricker, B.H., 2005. Therotocosis following the use of Ashwagandha. *Nederlands tijdschrift voor geneeskunde* 149, 2637–2638.
- van Grinsven, M., Parkipuny, L.M., Johns, T., 1999. *Euclea divinorum* (Ebenaceae) bark is a high-potential tanning material. *Economic Botany* 53, 220–221.
- Van Puyvelde L.M., Ngaboyisonga, P.C. and Rwangabo, S., 1977. Enquêtes ethnobotaniques sur la médecine traditionnelle rwandaise. Tome 1: Préfecture de Kibuye. Univ. Nat. Inst. Nat. Rech. Sc., Butare (Rwanda).
- van Rijn, C.P., Tanigoshi, K.L., 1999. The contribution of extrafloral nectar to survival and reproduction of the predatory mite *Iphiseius degenerans* on *Ricinus communis*. *Acarology Bulletin* 23, 281–296.
- Vasudevan, P., Kashyap, S., Sharma, S., 1997. Tagetes: a multipurpose plant. *Biore-source Technology* 62, 29–35.
- Vellaikumar, S., 1997. Comparative investigations on Chinaberry (*Melia azadirachta* L.) and Neem (*Azadirachta indica* A. Juss.) seeds as sources of insect antifeedants. MSc. Thesis, IARI, New Delhi, India.
- Ven Murthy, M.R., Ranjekar, P.K., Ramassamy, C., Deshpande, M., 2010. Scientific basis for the use of Indian ayurvedic medicinal plants in the treatment of

- neurodegenerative disorders: Ashwagandha. Central Nervous System Agents in Medicinal Chemistry 10, 238–246.
- Viable Herbal Solutions, 2003. About Olive Leaf-*Olea europaea* L. <http://www.viable-herbal.com/herbdesc3/1olive.htm> (27.10.03).
- Villamil, S.M., Alche, L.C.E., Coto, C.E., 1995. Inhibition of herpes simplex virus type-1 multiplication by meliacine, a peptide of plant origin. Antiviral Chemistry and Chemotherapy 6, 239–244.
- Volfová, A., Patočka, J., 2003. Strychnin-historie a současnost. Vojenské Zdravotnické Listy 72, 110–113.
- Wandibba, S., 1998. Who cares about the past? An example from Western Kenya. World Archaeological Congress: The Destruction and Preservation of Cultural Property, held from 3th to 7th May 1998. Island of Brac, Croatia, University of Southampton, Southampton.
- Wanzala, W., Sika, K.F.N., Gule, S., Hassanali, A., 2004. Attractive and repellent host odours guide ticks to their respective feeding sites. Chemoecology 14, 229–232.
- Warry, W., 1992. The eleventh thesis: applied anthropology as praxis. Human Organization 51, 155–163.
- Waters-Bayer, A., Bayer, W., 1994. Planning with Pastoralists: PRA and More. A Review of Methods Focused on Africa. Germany Agency for Technical Cooperation (GTZ), Eschborn, Germany.
- Watt, J.M., Breyer-Brandwijk, M.G., 1962. The Medicinal and Poisonous Plants of Southern and Eastern Africa, second ed. E. and S. Livingstone Ltd., London.
- Wehmeyer, A.S., 1966. The nutrient composition of some edible wild fruit found in the Transvaal. South Africa Medical Journal 40, 1102–1104.
- Were, G.S., 1967. A History of the Abaluyia of Western Kenya: c. 1500–1930. East African Publishing House, Nairobi, Kenya.
- Wilcox, E.V., McGeorge, Wm., 1912. Sisal and the Utilisation of Sisal Waste. Bulletin of Hawaii Agriculture and Experiment Station, Hawaii, pp. 24–29.
- William, N.S., 1999. Natural Products Drug Discovery. <http://chemistry.uah.edu/Faculty/setzer/natprod.html> (08.10.03).
- Williams, L.A.D., 1993. Adverse effects of *Artocarpus altilis* Park. and *Azadirachta indica* (A. Juss) on the reproductive physiology of the adult female tick, *Boophilus microplus* (Canest.). Invertebrate Reproduction and Development 23, 159–164.
- Wilson, A., Mansingh, A., 2002. Pesticidal potential of tropical plants-VI. Toxic actions of ethanol extract of selected plants on *Cylas formicarius elegantulus* Summer and *Boophilus microplus* Canestrinii. In: Mansingh, A., Young, E.D., Yee, H.T., Delgoda, R., Robinson, D.E., Morrison, Y.E., Lowe, I.H. (Eds.), Proceedings of the First International Symposium on Utilization of Natural Products in Developing Countries: Trends and Needs, held between 10th and 14th July, 2000 at the Natural Products Institute, University of the West Indies, Mona, Kingston 7, Jamaica, West Indies, pp. 240–246.
- Woodland, D.W., 1997. Contemporary Plant Systematics, second ed. Andrews University Press, Berrien Springs, Michigan.
- Working Paper FGR/18E, 2001. State of Forest Genetic Resources in Kenya. Forest Genetic Resources Working Papers. Food and Agriculture Organization of United Nation (FAO) Sub-Saharan Africa Forest Genetic Resources Programme of the International Plant Genetic Resources Institute (IPGRI/SAFORGEN), Danida Forest Seed Centre (DFSC) and International Centre for Research in groforestry (ICRAF), Forest Resources Division, FAO, Rome, Italy. <http://www.fao.org/DOCREP/004/AB396E/ab396e00.htm#Contents> (28.10.03).
- World Agroforestry Centre, 2003. Agroforestry Database. The *Ekebergia capensis*. <http://www.worldagroforestrycentre.org/Sites/TreeDBS/Aft/speciesinfo.cfm?SpID=720> (06.10.03).
- World Agroforestry Centre, 2003. The Prunus Net Database. <http://www.worldagroforestrycentre.org/Sites/TreeDBS/prunus/prunus.htm> (09.11.03).
- Wu, S.J., Chang, S.P., Lin, D.L., Wang, S.S., Hou, F.F., Ng, L.T., 2009. Supercritical carbon dioxide extract of *Physalis peruviana* induced cell cycle arrest and apoptosis in human lung cancer H661 cells. Food and Chemical Toxicology 47, 1132–1138.
- Wu, S.J., Ng, L.T., Huang, Y.M., Lin, D.L., Wang, S.S., Huang, S.N., Lin, C.C., 2005. Antioxidant activities of *Physalis peruviana*. Biological and Pharmaceutical Bulletin 28, 963–966.
- Wu, S.J., Tsai, J.Y., Chang, S.P., Lin, D.L., Wang, S.S., Huang, S.N., Ng, L.T., 2006. Supercritical carbon dioxide extract exhibits enhanced antioxidant and anti-inflammatory activities of *Physalis peruviana*. Journal of Ethnopharmacology 108, 407–413.
- Xu, Z.Q., Xu, F.C., 1993. Isolation and partial characterization of a protease from *Agave sisalana*. Acta Botanica Sinica 35, 171–178.
- Yen, C.Y., Chiu, C.C., Chang, F.R., Chen, J.Y., Hwang, C.C., Hseu, Y.C., Yang, H.L., Lee, A.Y., Tsai, M.T., Guo, Z.L., Cheng, Y.S., Liu, Y.C., Lan, Y.H., Chang, Y.C., Ko, Y.C., Chang, H.W., Wu, Y.C., 2010. Beta-hydroxywithanolide E from *Physalis peruviana* (golden berry) inhibits growth of human lung cancer cells through DNA damage, apoptosis and G2/M arrest. BMC Cancer 10, 46.
- Yumoto, T., Yamagiwa, J., Mwanza, N., Maruhashi, T., 1994. List of plant species identified in Kahuzi-Biega National Park, Zaire. Tropics 3, 295–308.
- Zakir, U.R., Ahmad, S., Qureshi, S., Atiq, U.R., Badar, Y., 1991. Toxicological studies of *Melia azedarach* L. (flowers and berries). Pakistan Journal of Pharmaceutical Sciences 4, 153–158.
- Zullo, M.A.T., Azzini, A., 1989. Steroidal sapogenins in sisal. Bragantia 48, 21–26.